



Persistent organic pollutants in the polar regions and the Tibetan Plateau: A review of current knowledge and future prospects

Xiaoping Wang^{a, b, c, *}, Chuanfei Wang^{a, b}, Tingting Zhu^{a, c}, Ping Gong^{a, b}, Jianjie Fu^d, Zhiyuan Cong^{a, b, c}

^a Key Laboratory of Tibetan Environment Changes and Land Surface Processes, Institute of Tibetan Plateau Research, Chinese Academy of Sciences (CAS), Beijing, 100101, China

^b CAS Center for Excellence in Tibetan Plateau Earth Sciences, Beijing, 100101, China

^c University of Chinese Academy of Sciences, Beijing, 100049, China

^d State Key Laboratory for Environmental Chemistry and Ecotoxicology, Chinese Academy of Sciences, Beijing, 100085, China

ARTICLE INFO

Article history:

Received 27 September 2018

Received in revised form

15 December 2018

Accepted 21 January 2019

Available online 7 February 2019

Keywords:

Persistent organic pollutants (POPs)

Polar regions

Spatiotemporal distribution

Bioconcentration

Climate change

ABSTRACT

Due to their low temperatures, the Arctic, Antarctic and Tibetan Plateau are known as the three polar regions of the Earth. As the most remote regions of the globe, the occurrence of persistent organic pollutants (POPs) in these polar regions arouses global concern. In this paper, we review the literatures on POPs involving these three polar regions. Overall, concentrations of POPs in the environment (air, water, soil and biota) have been extensively reported, with higher levels of dichlorodiphenyltrichloroethane (DDT) and hexachlorocyclohexane (HCH) detected on the Tibetan Plateau. The spatial distribution of POPs in air, water and soil in the three polar regions broadly reflects their distances away from source regions. Based on long-term data, decreasing trends have been observed for most “legacy POPs”. Observations of transport processes of POPs among multiple media have also been carried out, including air–water gas exchange, air–soil gas exchange, emissions from melting glaciers, bioaccumulations along food chains, and exposure risks. The impact of climate change on these processes possibly enhances the re-emission processes of POPs out of water, soil and glaciers, and reduces the bioaccumulation of POPs in food chains. Global POPs transport model have shown the Arctic receives a relatively small fraction of POPs, but that climate change will likely increase the total mass of all compounds in this polar region. Considering the impact of climate change on POPs is still unclear, long-term monitoring data and global/regional models are required, especially in the Antarctic and on the Tibetan Plateau, and the fate of POPs in all three polar regions needs to be comprehensively studied and compared to yield a better understanding of the mechanisms involved in the global cycling of POPs.

© 2019 Elsevier Ltd. All rights reserved.

1. Introduction

The Arctic and Antarctic were for a long time assumed to be virtually devoid of anthropogenic contaminants, since these polar regions are characterized by minimal prevalence of industrial and agricultural operations. However, anthropogenic contamination in the Arctic and Antarctic ecosystems was first documented in the 1960s (Risebrou et al., 1967; Sladen et al., 1966); and since then, there has been increasing evidence showing that contaminants can

reach the Arctic and Antarctic via atmospheric and oceanic transport from lower latitudes [Ahrens et al., 2010; Arctic Monitoring and Assessment Programme (AMAP) assessment report, 1998, 2004, 2009, 2015, 2016, 2017, 2018; Burkow and Kallenborn, 2000; Cabrerizo et al., 2017; Carrizo et al., 2017; Fellin et al., 1996; Gambaro et al., 2005; Gregor and Gummer, 1989; Jurado and Dachs, 2008; Kallenborn et al., 2013; Kallenborn et al., 1998; Li and Macdonald, 2005; Moeller et al., 2010; Montone et al., 2003; Nost et al., 2018; Oehme and Mano, 1984; Pozo et al., 2017; Uhl et al., 2012; Wang et al., 2017b; Wania and Dugani, 2003; Woehrschimmel et al., 2013; Wong et al., 2018].

Among the various anthropogenic contaminants, persistent organic pollutants (POPs) are relatively more toxic, persistent, bioaccumulative and prone to long-range atmospheric transport

* Corresponding author. Key Laboratory of Tibetan Environment Changes and Land Surface Processes, Institute of Tibetan Plateau Research, Chinese Academy of Sciences (CAS), Beijing, 100101, China.

E-mail address: wangxp@itpcas.ac.cn (X. Wang).

(Jones and de Voogt, 1999). All these characteristics promote their widespread distribution across the globe, including regions where they have never been used (Wania and Mackay, 1993; Wania and Mackay, 1995). Additionally, when POPs travel through the atmosphere, they tend to move in stages (a repeated process of warm volatilization and cold deposition), which is known as the “hopper” effect (Jurado and Dachs, 2008). This is the reason why POPs have been found in the cold Arctic and Antarctic regions (Jurado and Dachs, 2008). Owing to the effects of POPs (toxicity and bioaccumulation) on wildlife, pesticides were first found in fat tissues of polar bears and Antarctic penguins in the late 1960s (Risebrou et al., 1967; Sladen et al., 1966). In addition to the Arctic and Antarctic, however, researchers also began to search for evidence of airborne POPs in other cold ecosystems and mountain environments (Blais et al., 1998; Chen et al., 2008; Daly et al., 2007; Grimalt et al., 2001; Morrissey et al., 2005; Wania and Westgate, 2008).

The region that encompasses the Himalaya–Hindu Kush mountain range and the Tibetan Plateau is widely known as the Third Pole (TP), with an average elevation of 4700 m. The plateau's pivotal role is due almost entirely to its height, which makes it peculiarly cold for its latitude – colder than anywhere else and containing the largest reserve of fresh water outside the polar regions (Yao et al., 2013). Thus, the similarities between the TP and the two other polar regions are their cold temperatures, remoteness, and considerable water storage. Given the close proximity to South Asia, where POPs have been extensively applied, the TP encounters more pressure from contamination by POPs (Sheng et al., 2013; Wang et al., 2008a; Wang et al., 2010a). On the other hand, the climate of the TP is more complicated and its land surface (including forest, grassland, bare soil, lakes and glaciers) is more diverse than the Arctic and Antarctic, where ice is the dominant medium. These discrepancies make the transport and accumulation of POPs in the TP region different from the two other polar regions.

Regarding research on POPs in the two main polar regions and the TP, to date (as of 2017), based on ISI Web of Knowledge, more than 2000, 300 and 100 articles can be retrieved on the Arctic, Antarctic and TP, respectively. Among these articles, 66 reviews could be found for the Arctic, two for the Antarctic, but only one for the TP region. Systematic monitoring of POPs in the Arctic has been conducted for decades; detailed information with regard to the bioaccumulation of POPs in polar bears, the risk exposure of the Arctic ecosystem to POPs, the temporal trend of POPs in the air of the Arctic, and how climate change is influencing the cycling of POPs has been successfully reviewed (Braune et al., 1999; Fromberg et al., 1999; Houde et al., 2011; Lao et al., 2017; Ma et al., 2016; McKinney et al., 2015; Riget et al., 2016; UNEP/AMAP report, 2011; Vorkamp and Riget, 2014). This body of work provides clear insights into the current and future impacts of POPs on the Arctic environment. By comparison, the long-term monitoring of POPs in the Antarctic is scarce, with most research having focused on the accumulation of POPs in biota (Baek et al., 2011; Ellis et al., 2018; Kallenborn et al., 2013; Mangani et al., 2001; Wang et al., 2017b). Then, for the TP, although a relatively limited number of studies have been conducted, there is increasing interest from scientific communities and the general public on researching POPs in this region of the world.

Given the growth in research and the associated literature in this field, we first use a science mapping method to screen and summarize the hot topics in POP research for each polar region. Intellectual milestones and the evolutionary stages of major insights and understanding are extracted. Based on the outcome of this mapping process, the aims of the study are to: (1) summarize the main outcomes and common views for the main matrix of the ecosystem, including the air, water, soil and biota; (2) address the

influence of climate change on the transport and accumulation of POPs in the three polar regions, especially with respect to the release of chemicals by glacial melt; (3) synthesize the fate of POPs in the environments of the three polar regions; and (4) identify knowledge gaps and suggest where future studies should be directed.

Because AMAP – Arctic Council Working Group had published detailed assessment reports regarding fate, wildlife risk and human exposure of POPs in Arctic, this review would mainly focus on the data comparison and impact of climate change on POPs' accumulation at all three polar regions.

To the best of our knowledge, this is the first review that compiles the available achievements obtained for the two traditional polar regions, along with the TP, under the prohibition of POPs stemming from the Stockholm Convention, and provides assessments of the knowledge gaps in the scientific literature to promote best practice towards the highest quality research.

2. Summary of the hot topics in polar POP research

Pie charts (Fig. 1) were prepared based on the number of studies for different matrices over the two traditional polar regions and the TP. Research on the accumulation of POPs in biota is the dominant topic for the Arctic and Antarctic, followed air-, water-, sediment- and soil-related research. However, on the TP, POP research has focused mainly on atmospheric monitoring and transport mechanisms, with little reference to biota (Pan et al., 2014; Ren et al., 2017a; Wang et al., 2015a; Yang et al., 2011; Yang et al., 2010; Zhu et al., 2013). The temporal trends in the number of studies for the Arctic, Antarctica and TP are also shown in Fig. 1 (right-hand panel). After that the Stockholm Convention came into force in 2004 (<http://www.pops.int>), the international community got broadened and then commenced discussions on the need to eliminate or severely restrict POP production, thus, the amount of studies published with the aim of detecting and monitoring POPs in the Arctic and in Antarctica increased. In contrast, research on the TP started later, from 1997 (Galassi et al., 1997). This suggests that research interest has shifted from the traditional polar regions to a broader range that includes other remote areas.

Pie charts were also used to identify the important topics of wide concern among scientific researchers (Fig. 1). Regarding atmospheric POP research, the concentrations, spatiotemporal distribution and transport mechanisms (entering and moving out of the three polar regions) are the top three questions raising global concern. With respect to POPs in water, their concentrations in ice, snow, glacial runoff, remote lakes and ocean currents have been widely reported; plus, data from atmospheric monitoring and on the air–water (sea) exchange of POPs have attracted wide attention. Numerous publications have focused on the accumulation of POPs in biota, including the trophic dynamics of POPs throughout food webs, and the human exposure and health risk of POPs. Compared with research on air, water and biota, POP studies related to soils and sediments are rare, with those that have been conducted having focused mainly on the contamination of soil by POPs in the context of oil pollution in the Arctic and in Antarctica, and the related microbial diversities. However, a detailed soil survey has been conducted for the TP, in which both polluted areas in South TP and remote areas (hinterland of the Tibetan Plateau) were considered.

3. Atmosphere

Given the atmosphere is sensitive to emissions of pollutants, it reflects ongoing emissions, responds to source reductions, and acts as a carrier to disperse POPs globally (Bossi et al., 2013; Burkow and




Fig. 1. Publication statistics in different years for various media in the Arctic, Antarctic and Tibetan Plateau. (note: human biomonitoring studies were included in the biota class).

Kallenborn, 2000; Gregor and Gummer, 1989; Hung et al., 2001; Kallenborn et al., 1998; Kallenborn et al., 2012; Moeller et al., 2010; Oehme and Mano, 1984; Prevedouros et al., 2004; Wania and Mackay, 1993; Welch et al., 1991; Wong et al., 2018). Thus, atmospheric monitoring is essential for the three polar regions, as it can reflect the extent of POP reduction under the regulation of the Stockholm Convention and represent how emission source regions influence remote areas (Hung et al., 2016; Kallenborn et al., 2013; Wang et al., 2017a; Wang et al., 2017b; Wang et al., 2016d). In this section, the observation stations conducting long-term atmospheric monitoring programs are summarized; and the concentrations provided by these stations are compared to identify which remote parts of the Earth are facing the greatest pressure with

respect to POPs. The mechanisms and driving forces behind the atmospheric transport of POPs are then compiled.

3.1. Observation stations

There are five long-term observation stations located in the Arctic – namely, Alert [82°30'N, 62°20' W; 200 m above sea level (masl), on Ellesmere Island, Canada], station Nord (81°36'N, 16°40'W; 24 masl, on Greenland), Stórhöfði (63°24'N, 20°17'W; 118 masl, on Iceland), Zeppelin (78°54'N, 11°53'E; 474 masl, at Ny-Ålesund on Svalbard), and Pallas (68°00'N, 24°15'E; 340 masl, in Arctic Finland) (Fig. 2a). Arctic air monitoring research was established in the early 1990s and, to date, more than two decades of




Fig. 2. Research stations for monitoring atmospheric POPs in the (a) Arctic, (b) Antarctic and (c) Tibetan Plateau.

data have been accumulated, providing invaluable information on the spatial distribution and transport processes of POPs, which can be combined with various environmental transport and fate models to assess the long-range transport of POPs.

Compared with the Arctic, the maximum consecutive sampling time for the monitoring of POPs in the Antarctic atmosphere is three years, which includes the monitoring of atmospheric polychlorinated biphenyls (PCBs) and polybrominated diphenyl ethers (PBDEs) at the Chinese Great Wall Station on King George Island (Wang et al., 2017b, 2018), the monitoring of organochlorine pesticides (OCPs) and PCBs at King Sejong Antarctic station (Baek et al., 2011), and the monitoring of 32 PCB congeners, alpha- and gamma-hexachlorocyclohexane (HCH), trans- and cis-chlordane, trans- and cis-nonachlor, p,p'- and o,p'-DDT, dichlorodiphenyldichloroethane (DDD), dichlorodiphenyldichloroethylene (DDE), as well as hexachlorobenzene (HCB), at the Norwegian research station Troll (Dronning Maud Land) (Kallenborn et al., 2013). Besides this, atmospheric measurements of POPs have also been conducted at a monthly (Luek et al., 2017) and annual (Choi et al., 2008; Li et al., 2012a; Li et al., 2012b; Montone et al., 2003; Pozo et al., 2017) timescale, during multiple cruise-based expeditions and surveillance studies at Antarctic stations by various scientific teams. A detailed map of the sampling stations is presented in Fig. 2b.

Long-term air-monitoring observations (starting from 2008) have been conducted in the TP, but the stations are mainly located on the Tibetan Plateau, with very few observations available in North Pakistan and the Nepalese Himalaya (Sheng et al., 2013; Wang et al., 2017a; Wang et al., 2010a). Lulang is a station close to the Yarlung Tsangpo Grand Canyon (the entrance valley of the Indian monsoon), where relatively high OCP concentrations have therefore been observed (Sheng et al., 2013). Based on six years of monitoring, a recent study found a declining trend of OCPs in the atmosphere at Lulang (Wang et al., 2017a). Moreover, in 2007, a passive air-monitoring network comprising 16 sampling sites across the TP was established,

which provides data on the levels of POPs at a large spatial scale, offering insights into both the spatial and temporal trends of POPs (Wang et al., 2010a; Wang et al., 2016d). Fig. 2c summarizes all the long-term and short-term air-monitoring stations located on the Tibetan Plateau.

To our knowledge, the above-mentioned 5 Arctic stations, Chinese Great Wall Station in Antarctic and Lulang station in the TP had continued POPs monitoring to date. As remote areas of the world, long-term POP observations at these stations can provide accurate data on the temporal variations of POPs, and therefore connect the linkage between the temporal patterns of POPs and climate systems.

3.2. Comparison of atmospheric POP concentrations between the Arctic, Antarctic and TP

Since the 1990s, atmospheric monitoring of POPs has been conducted at four Arctic stations – Alert, Stórhöfði, Zeppelin, and Pallas – using traditional high-volume active air samplers (AASs) (Fellin et al., 1996). Therefore, 20 years of atmospheric data on legacy POPs and PCBs, as well as 10 years of data on PBDEs, have been obtained for this region (Hung et al., 2016). Apart from this comprehensive monitoring program, other relatively short-term observations have also reported various POP concentrations in the Arctic air, including some research using passive air samplers (PASs). Detailed concentration data for legacy POPs and emerging compounds, such as perfluorinated alkyl substances, EH-TBB (2-ethylhexyl-2,3,4,5-tetrabromobenzoate), HBB (hexabromobenzene), Dec 602 (2-ethylhexyl-2,3,4,5-tetrabromobenzoate), TCEP [Tris (2-chloroethyl) phosphate], TCPP [Tris (chloroisopropyl) phosphate], TPP (Tri-phenyl phosphate), TBEP [Tris (2-butoxyethyl) phosphate], and ΣOPEs (organophosphate esters), are presented in Supporting Information (SI) Table SI-1. Similarly, the AAS data obtained from the Chinese Great Wall Station in Antarctic and other PAS data are reported in Table SI-2. The compounds include legacy POPs and PBDEs (Table SI-2).

To date, the AAS data have been gathered on the Tibetan Plateau at the stations of Lulang, Namco, Lhasa, Mt. Everest, Mt. Yulong, Nagri, and Waliguan; while PAS data cover 16 sites across the plateau (Wang et al., 2016c). Table SI-3 compiles all the reported concentrations for both legacy POPs and emerging compounds [TBE: 1,2-bis(2,4,6-tribromophenoxy)ethane, DBDPE: decabromodiphenyl ethane, syn-DP: syn-dechlorane plus, anti-DP: anti-dechlorane plus, DP: dechlorane plus].

In order to compare the differences in concentrations of chemicals between the Arctic, Antarctic and the Tibetan Plateau, a bar graph showing the minimum and maximum concentrations of chemicals was prepared (Fig. 3). The results reveal the following general findings: (1) the concentrations of chemicals in the Antarctic are generally lower than those of the Arctic; (2) similar concentration ranges of HCB, α -HCH, γ -HCH and α -endosulfan are apparent between the Arctic and Tibetan Plateau; (3) wide ranges of PCB concentrations occur in the Antarctic, spanning six orders of magnitude, while the highest PCB concentrations for the three polar regions are similar, within a factor of three; (4) different from other chemicals, PBDE concentrations on the Tibetan Plateau are obviously lower than those for the Arctic and Antarctic; (5) interestingly, short-chain chlorinated paraffin (SCCP) concentrations on the Tibetan Plateau are one and two orders of magnitude higher than those of the Arctic and Antarctic, respectively (Fig. 3). Fig. 3 gives a direct insight into the levels of chemicals over the three polar regions.

3.3. Transport mechanisms

Contaminants are transported to the Arctic, Antarctic and TP by




Fig. 3. Comparison of the concentration ranges of atmospheric POPs in the Antarctic, Arctic and Tibetan Plateau. Data and references used in this figure can be found in Table SI 1–3.

various pathways (Becker et al., 2011; Benskin et al., 2012b; Carrizo et al., 2017; Gong et al., 2014; Gong et al., 2015; Kallenborn et al., 2007; Kallenborn et al., 2012; Li et al., 2002; Liu et al., 2010; Moeller et al., 2010; Sheng et al., 2013; Sobek and Gustafsson, 2004), but the primary pathway is atmospheric transport. Many POPs are transported in the atmosphere to the Arctic from densely populated areas at lower latitudes, with subsequent deposition and revolatilization (Burkow and Kallenborn, 2000; Wania et al., 2017; Wania and Mackay, 1995; Wania and Mackay, 1999; Wania et al., 1999). This has been termed “the grasshopper effect” and explains how POPs travel extremely long distances and arrive in remote and cold areas (Gouin et al., 2004; Jurado and Dachs, 2008). Atmospheric transport is easily influenced by seasonal weather-related fluctuations. For example, during winter, high-pressure systems situated in Siberia push air masses north into the Arctic, which accounts for much of the Eurasian atmospheric contaminant input to the Arctic (Halsall et al., 1997). As presented by UNEP/AMAP report (2011), the climatic impact on the atmospheric transport of POPs has been investigated. ENSO (El Niño–Southern Oscillation) plays an important role in the atmospheric concentration of POPs in the Arctic. Besides, fluctuations of the AO (Arctic Oscillation) and NAO (North Atlantic Oscillation) can also change strength of the air pressures and then vary the directions of westerly winds in Arctic and Atlantic, and further change the levels and transport of POPs (Ma et al., 2004; Macleod et al., 2005; Octaviani et al., 2015). However, owing to the data on POP concentrations in the Antarctic atmosphere being relatively short term, they have yet to be linked with atmospheric circulation patterns (Kallenborn et al., 2013; Wang et al., 2017b). To date, for the TP, there are only six years of AAS POP observations available, at Lulang in the southeast of the region. Results showing almost the same seasonal fluctuation of POPs with the Indian Monsoon Index indicate that the Indian Monsoon wind drives the transport of POPs, pluming through the South Asian continent and importing POPs from there onto the Tibetan Plateau (Sheng et al., 2013; Wang et al., 2017a). Climate can influence the spatial and seasonal patterns of POPs by changing the air circulation/transport paths.

4. Vegetation

POPs can be scavenged from the atmosphere by plant leaves, the vegetation in Arctic, Antarctic and the TP is an important receptor of airborne POPs. Lichens and mosses are dominant species in Polar regions and had been considered good indicators of atmospheric

pollution (Cabrerizo et al., 2012; Sanchis et al., 2015; Yogui and Sericano, 2008; Yogui et al., 2011). Banned legacy POPs, such as HCB, DDE, PCBs (Cabrerizo et al., 2012) (Yang et al., 2013) and other ongoing pollutants, such as, perfluorinated alkyl substances (PFASs, Moeller et al., 2011), short-chain chlorinated paraffins (SCCPs, Lao et al., 2017), dechloranes (Yang et al., 2016) and volatile dimethylsiloxanes (VMS) (Sanchis et al., 2015) were measured in different vegetation types at three Polar regions (Augusto et al., 2013; Yang et al., 2013; Zhu et al., 2015b). Generally, lipid and wax contents of vegetation strongly affect the accumulation process, moreover, leaf morphology, physiological properties, growth rates are also responsible factors for accumulating organic air pollutants in vegetation (Franzaring and van der Eerden, 2000). Plant species with a high surface-to-volume ratio usually accumulate more organic air pollutants than species with compact leaves (Franzaring and van der Eerden, 2000). Given the partitioning of organic pollutants from air to vegetation is controlled by ambient temperatures, with global warming, volatile compounds may return to the gas-phase (Barber et al., 2004). Another contrary situation is that enhanced vegetation biomass by warm temperature will provide more sorption surface for POPs' accumulation (Cabrerizo et al., 2013).

5. Ice and snow

Due to its porous structure and low temperatures, snow can effectively scavenge both gas phase and particle-bound chemicals from the air and thus mainly contribute to the atmospheric deposition of POPs in three Polar Regions. The deposition of snow POPs can be transferred deeper into aged snowpack and glacier to record the global scale long range atmospheric transport for a long time. Up to date, legacy pesticides, current use pesticides, emerging PFASs and brominated flame retardants (BFRs) had been widely detected in ice capes of Arctic [most of samplings were conducted at Svalbard, Norway (Garmash et al., 2013; Hermanson et al., 2010; Hermanson et al., 2005; Kwok et al., 2013; Ruggirello et al., 2010) and Devon Ice Cap (Meyer et al., 2012; Young et al., 2007; Zhang et al., 2013), Canada] and Antarctic [King George Island (Cipro et al., 2017) and Talos Dome (Fuoco et al., 2012)]. Because the mountain glacier at high altitude usually have limited snow melt, the deposition history of POPs can be relatively stored at high temporal resolution. In the TP, DDT and HCH records were retrieved from Mt. Everest (Wang et al., 2008a) and Dasuopu glacier (Wang et al., 2008b); PFASs temporal trends were observed from Zuoqiupu and Muztagata glacier (Wang et al., 2014a,b,c,d).

Table 1
Concentrations of POPs in ice and snow from the Arctic.

Unit	Sampling time	α -HCH	γ -HCH	Σ DDTs	HCB	Σ PCBs	Σ PFASs	Ref.
		pg/L	pg/L	pg/L	pg/L	pg/L	pg/L	
Canadian Arctic	ice 1993					1200–6700 (3500) ^a		Gregor et al. (1995) Peters et al. (1995) Pucko et al. (2010)
	ice 1993							
	ice 2007–2008	4.013 \pm 0.307	0.423 \pm 0.013					
	snow 2010						260–1500	Cai et al. (2012b)
	ice 2010						2200	Cai et al. (2012b)
Central Arctic	snow 2001					2.4–6.9 ^b	134–848	Veillette et al. (2012)
	ice 2001					2–15 ^c		Gustafsson et al. (2005) Gustafsson et al. (2005)
	ice 2006							(Kwok et al., 2013)
Norwegian Arctic	snow 2003	17–382(130)	265–4390(1500)	1.73–80.5(18.8)	9.51–62.5(22.8)	160–2540 (1060) ^d	80–200	Herbert et al. (2005)
	snow 2015	34.7–48.2						Bigot et al. (2017)
North Greenland	snow 2015							

^a Σ_{67} PCBs.

^b Σ_{15} PCBs.

^c Σ_{15} PCBs.

^d Σ_{14} PCBs.

POPs concentration in snow/ice of polar regions (Table 1) and the TP (Table SI-4) were compared, displaying concentrations of HCB and DDTs in glaciers in the TP are higher than those in the Arctic; meanwhile, PCBs are lower than in the Canadian Arctic and Norwegian Arctic (Table 1 and Table SI-4). Concentration of PFASs in snow/ice cores broadly followed the order of Tibetan fresh snow > Tibetan ice core > Arctic glacier > ice cap of Antarctic (Kwok et al., 2013; Wang et al., 2014a,b,c,d; Young et al., 2007).

Temporary trends of PBDEs (Hermanson et al., 2010; Meyer et al., 2012), PCBs (Fuoco et al., 2012; Garmash et al., 2013; Zhu et al., 2015a), DDTs (Wang et al., 2008a), HCHs (Wang et al., 2008a) and PFASs (Wang et al., 2014a,b,c,d) (Young et al., 2007; Zhang et al., 2013) had been retrieved from ice cores. These trends were overall the representative of emission history of chemicals, however, due to the relocation of chemicals caused by post-depositional processes, the retrieved atmospheric deposition history exhibited some uncertainties (Pavlova et al., 2014; Zhang et al., 2013). For example, in Arctic, three PCB flux peaks were observed from 1957 to 1966, in 1974–1983, and 1998–2009 (Garmash et al., 2013). While, in Antarctic, sharp concentration increase occurred in 1930–1980, but a constant trends was from 1980 to 2002 (Fuoco et al., 2012). Both emission patterns and melting condition can lead to the above difference. An interesting phenomenon was that PCB-11 was homogeneously found in Arctic and Antarctic ice and snow (Fuoco et al., 2012; Garmash et al., 2013; Vecchiato et al., 2015), although it is not a commercial PCB product. Incineration of paper and plastics containing diarylide pigments were suggested as a major source of PCB-11 toward the Arctic (Vecchiato et al., 2015), however, its source for Antarctic is not clear.

With global warming, POPs can enter the environment of the polar regions, via the melting of glaciers and sea ice, has come to the fore (Bigot et al., 2017; Bizzotto et al., 2009; Blais et al., 2001; Guzzella et al., 2016; Meyer et al., 2009; Zhao et al., 2012; Pavlova et al., 2014). In the Arctic, high concentration episodes of β -HCH were observed throughout the year at Alert, when the Arctic sea-ice extent reached a record low. The mechanism is believed to involve re-emissions from the oceans because of sea-ice retreat sustaining levels in the Arctic air (Ma et al., 2011). Furthermore, Kwok et al. (2013) investigated the transport of PFAS from an arctic glacier to downstream locations and found melting glacier is the secondary source of PFASs. Melting glaciers can also increase the concentration of POPs in Himalayan lake water (Sharma et al., 2015). Furthermore, under global warming, POPs can migrate into lakes through the water cycle, and even accumulate in lake sediments (Schmid et al., 2011).

6. Water

Besides the air, water has become another widely used environmental matrix for monitoring POPs (Braune et al., 1999; Fernandez et al., 2005; Muir and Lohmann, 2013; Yamashita et al., 2005; Yao et al., 2002). Different from in the air, acted on by temperature, pollutants in water will migrate with phase transition (i.e., gas, liquid and solid) (Bidleman, 1999; Bigot et al., 2017; Blais et al., 2001; Cai et al., 2012a). Remote water and glaciers have been seen as “sinks” of pollutants (Gregor and Gummer, 1989; Gregor et al., 1995; Kang et al., 2009; Peters et al., 1995; Wang et al., 2014c; Wang et al., 2008b). However, in response to climate warming, oceans and glaciers possibly re-release pollutants into the environment (Jantunen et al., 2015; Ma et al., 2016; Ma et al., 2011; McKinney et al., 2015; Riget et al., 2013). In this section, the levels and spatial distribution of POPs in water in the three polar regions are compared. Furthermore, the concentrations of water pollutants reaching the polar regions are summarized, and the connections between the air–water exchange of POPs and climate warming are presented.

6.1. Concentrations

Statistically, research on POPs in ocean water for the polar regions has mainly focused on the Arctic, with twice as much work having been carried out than on the Antarctic, and ten times more than that for the TP. The concentrations of POPs in ocean water in the polar regions are summarized in Table 2 for the Arctic, and Tables SI-4 and 5 for the TP and Antarctic respectively. In general, data are still limited for the TP, not only in terms of sampling sites but also with respect to the variety of species of pollutants (Chakraborty et al., 2016; Li et al., 2017b; Ren et al., 2017b). HCHs, DDTs, and PCBs are commonly concentrated pollutants in the three polar regions (Table 1). In the Arctic Ocean, concentrations of HCHs and PCBs show wide ranges, from 0.1 to 3.2×10^3 pg/L, and from 0.13 to 4.0×10^5 pg/L, respectively, and DDTs are mainly lower than 1 pg/L (Table 2). Concentrations of HCHs and DDTs in the TP are higher than those in the Arctic and Antarctic (Table 2). Concentrations of PCBs are highest in the Arctic, followed by the Antarctic and then the TP (Table 2).

For other pollutants, the average concentration of HCB in ocean water of the Arctic is higher than that of the Antarctic (Table 2). Endosulfan and PBDEs have only been reported in water in the Arctic, with concentrations in the ranges of ND–55 and 0.3–11.2 pg/L respectively (Table 2). The concentration of the emerging

Table 2

Concentrations of POPs in ocean water from the Arctic.

Unit	Sampling time	α -HCH	γ -HCH	Σ HCHs	Σ DDTs	HCB	Σ PCBs	Σ PBDEs	Endosulfan	Σ PFASs	OPEs	Ref.
		pg/L	pg/L	pg/L	pg/L	pg/L	pg/L	pg/L	pg/L	pg/L		
Arctic Ocean	surface water	1993	250						3.8			(1)
	deep water	2001, 2005, 2008					0.13–21 ^a					(2)
	deep water	2001, 2005, 2008						0.3–11.2 ^g				(3)
	deep water	2001, 2005, 2008			0.10–66							(4)
	surface water	2004					<1 ^b					(5)
	surface water	2008–2009		262–3156	ND-0.31				0.04–0.18			(6)
	surface water	2008	630–2690 (1978)	170–700 (382)					ND-55			(7)
	surface water	2008	0.12–0.76 (0.44)	0.20–0.37 (0.26)	0.1–5.14 (1.15)	ND-0.16 (0.09)			ND-0.1 (0.05)			(8)
	surface water	2010										(9)
Canadian Arctic	2 m	1998	1100 ± 7 00	200 ± 10	1400 ± 100							(10)
	7 m	2005–2009										(11)
	7 m	2007–2008	465–1013	150–254		4.0–6.4						(12)
	7 m	2007, 2008, 2010							0.20–2.3			(13)
	7 m	2010–2011								27–754		(14)
	7 m	2010–2011									DL-4400	(15)
Greenland	11 m	2009								1900–153000		(16)
	11 m	2009								14.5–449		(17)
	5 m	2010	11.00			50.00				50–280		(18)
	5 m	2015	224–253									(19)
Europe Arctic	8 m	2004	ND-65 (13)	ND-21 (4.7)	1.5–83 (16.9)	0.17–0.96 (0.4)	0.8–9.6 (4.6)					(20)
	8 m	2009						0.03–0.64 ^h				(21)
Norwegian Arctic	surface water	2009					2000–400000 ^c					(22)
	surface water	2009					11.6–27.7 ^d					(23)
Russian Arctic	5 m	2003–2005			291	3	9	12 ^e	5 ⁱ			(24)
	surface water	2010								500 ± 170		(25)
Western Arctic	8 m	1996	910	270								(26)
	8 m	2001					0.13–3.92 (1.34) ^f					(27)
	surface water	2012								10–1700		(28)
Bering Sea and Chukchi Sea		1999		226–652								(29)

(1)(Weber et al., 2006); (2)(Carrizo and Gustafsson, 2011a,b); (3)(Salvado et al., 2016); (4)(Carrizo et al., 2017); (5)(Gioia et al., 2008); (6)(Qiu and Cai, 2010); (7)(Jantunen and Bidleman, 1998); (8)(Cai et al., 2010); (9)(Ke et al., 2017); (10)(Moisey et al., 2001); (11)(Benskin et al., 2012b); (12)(Wong et al., 2011); (13)(Morris et al., 2016); (14)(Veillette et al., 2012); (15)(McDonough et al., 2018); (16)(Lescord et al., 2015); (17)(Busch et al., 2010); (18)(Zhao et al., 2012); (19)(Carlsson et al., 2012); (20)(Bigot et al., 2017); (21)(Lohmann et al., 2009); (22)(Moeller et al., 2011); (23)(Polkowska et al., 2011); (24)(Papale et al., 2017); (25)(Carroll et al., 2008); (26)(Cai et al., 2012b); (27)(Harner et al., 1999); (28)(Sobek and Gustafsson, 2004); (29)(Yeung et al., 2017); (30)(Yao et al., 2003).

^a Σ_{13} PCBs.^b Σ_6 PCBs.^c Σ_7 PCBs.^d Σ_9 PCBs.^e Σ_{10} PCBs.^f Σ_{12} PCBs.^g Σ_{14} PBDEs.^h Σ_{10} PBDEs.ⁱ Σ_4 PBDEs.

group of compounds, PFASs, in Arctic water, is comparable to that of King George Island but higher than that of the western Antarctic (Table 2).

6.2. Spatial distribution

Owing to studies on POPs in water in the Antarctic and TP being quite scattered, examining the spatial distribution of POPs in water has instead focused mainly on the Arctic. The Arctic Ocean is composed of the central Arctic Ocean and seven vast continental shelf seas (including the Chukchi, East Siberian, Laptev, Kara, Barents, Beaufort and Lincoln seas) that cover 53% of its waters (Figure SI-1) (Jakobsson, 2002). In the Arctic, large-scale observations have also been performed along the cruise tracks of Arctic Ocean expeditions (Benskin et al., 2012a; Cai et al., 2010; Cai et al., 2012b; Jantunen and Bidleman, 1998; Lohmann et al., 2009; Moeller et al., 2010; Sobek and Gustafsson, 2004; Xie et al., 2011; Yao et al., 2002). Overall, the concentrations of POPs in surface water have shown decreasing trends with increasing latitude, which is consistent with distances away from source regions (Benskin et al., 2012b; Carrizo et al., 2017; Gioia et al., 2008; Salvado et al., 2016; Sobek and Gustafsson, 2004; Yao et al., 2002). Longitudinally, the values of POPs tend to decrease from west to east, but this trend is not statistically significant (Yao et al., 2002). For different parts of the Arctic Ocean, the concentrations of POPs in the water of shelf-sea areas are usually higher than in the central Arctic ocean (Carrizo and Gustafsson, 2011a,b; Carrizo et al., 2017; Salvado et al., 2016; Sobek and Gustafsson, 2004).

Dissolved contaminants may reach deeper water layers through vertical mixing or convective overturning in well-mixed shelf areas (Carrizo et al., 2017; Sobek and Gustafsson, 2014a). The vertical profile of dissolved-phase POPs in surface seawater shows a general decreasing trend in concentration with depth, such as for DDT, HCB, Endosulfan, and the low-molecular-weight PCBs, and PBDEs, in the Arctic Ocean (Cai et al., 2010; Fernandez et al., 2005; Salvado et al., 2016; Sobek and Gustafsson, 2014a; Sun et al., 2016; Yamashita et al., 2005; Yamashita et al., 2008). However, in deep water, increasing concentrations of DDE and high-molecular-weight compounds with depth have been observed (Carrizo et al., 2017; Salvado et al., 2016; Sobek and Gustafsson, 2014b). For the vertical distribution of emerging compounds, the detection of PFAS has been shown to be limited to the polar mixed layer (15–30 m) and halocline (150 m below the surface); whereas, in the North Barents Sea, perfluorooctane sulfonate (PFOS) has been detected down to 250 m below surface (Yeung et al., 2017).

6.3. Oceanic and riverine transport

Aside from atmospheric transport, both water-soluble and particle-bound compounds can be transported via rivers or ocean currents (Alexeeva et al., 2001; Carroll et al., 2008; Sun et al., 2016). For example, HCH is generally regarded as a “swimmer”, being stored in rivers and ocean water; plus, owing to the “wash-out” of rivers, large areas of land can be flushed into the Arctic, providing an extra release of POPs into this region (Alexeeva et al., 2001; Carlsson et al., 2012; Carrizo and Gustafsson, 2011a,b; Carroll et al., 2008; Lohmann and Belkin, 2014; Nikanorov et al., 2015; Zhulidov et al., 2000; Zhulidov et al., 2002). Along with transport via oceanic currents, phytoplankton can also carry POPs into the oceans (Dachs et al., 2002; Galban-Malagon et al., 2012) – a mechanism referred to as the “biological pump”. In the Greenland current and Atlantic sector, the transport of PCBs and HCHs to the Arctic Ocean is strongly retarded by the oceanic biological pump (Galban-Malagon et al., 2012; Galban-Malagon et al., 2013). High settling fluxes of

PCBs due to the biological pump have also been observed in the Southern Ocean (Galban-Malagon et al., 2018).

6.4. Air–water exchange and its relationship with climate change

Gaseous exchange occurs constantly for POPs at the air–water interface (Bidleman, 1999; Jantunen and Bidleman, 1996; Jantunen et al., 2008; Wong et al., 2011). However, air–water gas exchange is a two-way process, which can alternate between net deposition and net volatilization in response to seasonally changing temperatures, atmospheric levels, cycles of biological productivity, and so on. Currently, the seasonal variations of air–water exchange are observed in three polar regions, but long-term trends are only available for the Arctic. In summer, HCB, γ -HCH and β -HCH are close to equilibrium or undergoing net deposition, with occasional reports of volatilization in some areas or seasonal variations in the polar oceans and lake water of the TP (Bidleman et al., 2007; Bigot et al., 2016; Cai et al., 2012a; Cincinelli et al., 2009; Ding et al., 2007; Ren et al., 2017b). Currently used pesticides, such as α -endosulfan, are generally dominated by net deposition (Weber et al., 2006).

Based on long-term data in the Arctic, the connection between air–water exchange and climate change has been explored using both observational data and models (Jantunen et al., 2015; Ma et al., 2016; Ma et al., 2011; McKinney et al., 2015; Xie et al., 2011). The direction of α -HCH shifted from net deposition in the 1980s to net volatilization in the 1990s (Fig. 4), which was influenced by declining primary emissions and the climate warming (Cai et al., 2012a; Ding et al., 2007; Jantunen and Bidleman, 1996; Jantunen and Bidleman, 1995; Jantunen et al., 2008). Since 1990, the net direction of α -HCH gaseous exchange has likely been controlled more by global warming and rapid changes in sea-ice coverage than by changes in global emissions of α -HCH (Cai et al., 2012a; Wong et al., 2011). Based on an atmospheric model for POPs, Ma et al. (2011) they found that other toxic chemicals (such as p,p'-DDT, PCB congeners 28, 52 and 101) probably volatilize from secondary emission sources/reservoirs in water, snow and ice across the Arctic under climate warming (Ma et al., 2011). Therefore, under Arctic warming, some POPs in water will return back to the atmosphere, becoming available for circulation once again, which will in turn offset the effect of reducing the primary emission sources.




Fig. 4. Temporal trends for air–water gas exchange fugacity ratios and flux of α -HCH in the Arctic Ocean [data are from Cai et al. (2012a), Ding et al. (2007), Jantunen and Bidleman (1996), Jantunen and Bidleman (1995) and Jantunen et al. (2008)].

7. Soil and sediment

Soil and sediment in the cold polar regions and high mountains provide stationary “end points” where POPs possibly reside for the long term (Borghini et al., 2005b; Chen et al., 2008; Wania and Mackay, 1993; Wania and Westgate, 2008). Without obvious local emission sources, soil and sediment in the polar regions act as natural recorders of POP levels over long timescales, thus accumulating valuable information on past environmental and climatic events.

7.1. Soil

Comparisons of POPs in the soils of the three polar regions are summarized in Table 3, from which it is clear that the levels/orders of magnitude are similar to those mentioned above for water. The spatial distribution of POPs in soil in the Arctic and Antarctic has mainly been examined on small scales (specific islands) (Borghini et al., 2005a; Cabrerizo et al., 2013; Casal et al., 2018; Klanova et al., 2008; Negoita et al., 2003; Park et al., 2010; Wang et al., 2012a; Wang et al., 2015b). Whereas, studies on POPs in soil in the TP have been carried out across the plateau on a larger scale (Tao et al., 2011; Wang et al., 2014a; Wang et al., 2012c; Zheng et al., 2012). Generally, concentrations show decreasing trends from the edge to the hinterland areas of the TP, and the profile is dominated by more volatile compounds (Klanova et al., 2008; Park et al., 2010; Wang et al., 2012a; Wang et al., 2012b; Wang et al., 2014a; Wang et al., 2012c), which reflects well the long-range atmospheric transport of POPs. Besides, higher levels of POPs have been observed sporadically near research stations (Abakumov et al., 2014; Aislabie et al., 1999; Li et al., 2017a; Park et al., 2010; Wang et al., 2014a; Wang et al., 2016b) and military bases (Bright et al., 1995; Stow et al., 2005) in the polar regions or populated regions of the TP, indicating local sources at these sites.

Soil in cold regions are considered as “sinks” of POPs (Meijer et al., 2003), but the direction of air–soil gas exchange of POPs may change with variations in temperature and soil organic matter (SOM) (Wang et al., 2012c). Cabrerizo et al. (2013) provided strong evidence that both climate change (temperature) and carbon cycling (SOM content) drive the remobilization of PCBs in the Antarctic, not only due to warming. In the TP, grassland and forest soil are sinks of DDTs, while desert soil might re-release low-

molecular-weight PCBs, induced by variations in SOM (Wang et al., 2014b; Wang et al., 2014d; Wang et al., 2012c).

7.2. Sediment

Sediments are recognized as a useful record of atmospheric deposition (Fernandez et al., 2000; Malmquist et al., 2003; Martins et al., 2010; Rawn et al., 2001; Sun et al., 2005; Zaborska et al., 2011). The geographical distributions of POPs in surface sediment are usually similar to those for soil or water; plus, the temporal variations of POPs vertically in sediments provide a record of their emissions history globally (Cheng et al., 2014a; Cheng et al., 2014b; Combi et al., 2017; Dong et al., 2015; Duan et al., 2011; Galassi et al., 1997; Guo et al., 2011; Jin et al., 2017; Jin et al., 2015; Li et al., 2017b; Lin et al., 2017; Ma et al., 2017a; Ma et al., 2017b; Malmquist et al., 2003; Sobek and Gustafsson, 2014a; Sun et al., 2018; van Drooge et al., 2013; Wang et al., 2010b; Yang et al., 2016). Both of these aspects have been extensively reported and will not be reviewed here. Instead, in this section, the shifts in the sources of POPs in sediments over long timescales, along with the relationships between POP levels and sedimentary characteristics, are the focus.

Studying the temporal variation in the concentrations and relative compositions of POPs has generally been the main approach to tracing the possible sources of such pollutants (Blais et al., 2001; Dahle et al., 2003). Against the background of climate warming, glacial melt has become another dominant source of POPs for nearshore waters adjacent to glaciers and lakes (Bigot et al., 2017; Blais et al., 2001; Cai et al., 2012b; Sharma et al., 2015). In the sediments of glacial lakes, a second peak of DDTs and HCHs was observed in the 1990s after the ban of these compounds (Cheng et al., 2014a; Li et al., 2017b; Sun et al., 2018) – a phenomenon believed to derive from the melting of glaciers (Blais et al., 2001).

Besides sources, sedimentary properties [including grain size, water content, loss on ignition, total organic content (TOC), and black carbon] might play a role in the vertical distribution of POPs (Fernandez et al., 1999; Hong et al., 2012; Li et al., 2017d; Ma et al., 2015; Xue et al., 2016). Thus, the relationships between POPs and sedimentary properties have been explored in several studies. For example, no apparent co-relationships were obtained between PCB concentrations and the abovementioned sedimentary properties for the Bering Sea, Chukchi Sea, and Canada Basin (Hong et al.,

Table 3
Comparison of POPs in soils from the polar regions and the Tibetan Plateau (TP).

Sampling sites	Sampling time	Σ HCHs pg/g	Σ DDTs pg/g	HCB pg/g	Σ PCBs pg/g	Σ PBDEs pg/g	Ref.
Arctic							
Ny-Ålesund	2013					42	Wang et al. (2015b)
Ny-Ålesund	2007						Wang et al. (2009b)
Island of Tromsø	2013–2014	NA-81		616–1477	300–950		Casal et al., (2018)
Antarctic							
Livingston Island	2009				24.5–40.9		Cabrerizo et al., (2013)
Eastern coast of Antarctica	1998	86–469	110–1220	20–25280	200–410		Negoita et al., (2003)
King George Island	2006				8.0–33.8		Park et al., (2010)
Tibetan Plateau							
Tibet		670–5380	ND-2830				Fu et al., (2001)
Mt. Qomolangma	2005				47.1–422.6	4.3–34.9	Wang et al. (2009a)
Tibetan plateau	2005–2006	467 ± 741	329 ± 818				Tao et al., (2011)
	2007	64–847	13–7700	24–564	75–1021	ND-27	Wang et al. (2012c)
	2008				163	26	Zheng et al., (2012)
	2011						Wang et al., (2013)
	2007						Wang et al. (2014a)
	2014	430–5690	21–1230				Tao et al., (2014)
Maizhokunggar County	2015						Li et al. (2017c)
Western TP	2014						Zhou et al., (2018)

2012), and grain size was found to be a minor factor for the accumulation of POPs (Li et al., 2017d; Xue et al., 2016). However, significant relationships of PBDEs with total organic carbon (TOC) and black carbon have been found (Cai et al., 2012c; Xue et al., 2016). These observations indicate that, besides sources, the distribution of POPs in sediment is controlled by multiple factors, such as atmospheric transport and deposition, mixing, partitioning, and sorption in the water column and sediments.

8. Biota

Due to lipophilicity, POPs can bioaccumulate in organisms and are stored in the lipid component of tissues (Muir et al., 1988). The bioaccumulation and biomagnification of POPs along aquatic and terrestrial food chains have been widely reported (Corsolini et al., 2017; Corsolini et al., 2003; Corsolini et al., 2002; Corsolini and Sara, 2017; Evenset et al., 2016; Goerke et al., 2004; Hop et al., 2002; Kelly and Gobas, 2001; Kelly and Gobas, 2003; Lescord et al., 2015; Morris et al., 2016). Because of their distance from emission sources, biota in the polar regions can serve as adequate biological indicators of global POP emissions (Braune et al., 1999; Braune, 2007; Braune et al., 2002; Braune et al., 2015; Braune and Mallory, 2017; Helgason et al., 2008; Riget et al., 2010a; Riget et al., 2013; Riget et al., 2010b). To date, the levels, bio-concentration mechanisms, and temporal trends of POPs in biota from the polar regions have been extensively reported and reviewed by other papers (Braune et al., 1999; Cannon et al., 2016; Corsolini, 2009; Houde et al., 2011; Letcher et al., 2010; McKinney et al., 2015; Vorkamp, 2016) and AMAP assessment reports (2009, 2015, 2018). Here, the factors influencing the levels and bio-accumulation of POPs in biota in the polar regions, as well as their long-term trends and exposure effects, are reviewed.

8.1. Factors influencing levels and bioaccumulations

The levels of POPs in the Arctic, Antarctic and TP vary by region, biota type, and even tissue type (Braune et al., 1999; Braune et al., 2002; Braune et al., 2005; Corsolini et al., 2003; Court et al., 1997; Inomata et al., 1996). For example, differences have been observed between benthic and pelagic species (Goutte et al., 2013; van den Brink et al., 2011). Goutte et al. (2013) found levels of HCB to be far higher in pelagic than in benthic organisms of Antarctic. Also, concentrations in marine biota have been found to be higher than those in freshwater and terrestrial ecosystems (Dietz et al., 2000). Among the different tissues, liver tissue, with its relatively higher lipid content, accumulates higher concentrations of POPs (Aas et al., 2014; Andersson et al., 1998).

Biological (lipid dynamics, food-web productivity and migration) and environmental (emissions and climate change) factors possibly play a role in bioaccumulations of POPs in the polar regions (Andersson et al., 1998; De Laender et al., 2010; Fisk et al., 2001; Gustayson et al., 2015; Li et al., 2016; Pedro et al., 2017; Rios et al., 2017). In response to the limited observational data available in the polar regions, models have been set up to examine to what extent these factors cause bioaccumulation in biota (Borga et al., 2010; De Laender et al., 2010; Kelly and Gobas, 2003). For example, driven by lipid dynamics, “bioaccumulation factors” in fish were found to be 10 times lower in summer than in spring and fall/winter (De Laender et al., 2010); and driven by seasonal shifts in trophic levels and lipid dynamics, “trophic magnification factors” were highest for PCB 153 during spring and lowest for PCB 52 in summer and fall/winter (De Laender et al., 2010). Furthermore, increased temperatures and emissions were found to reduce the overall bioaccumulation of organic contaminants in the Arctic marine food web, with the largest change being for PCB-52 and PCB-153 (Borga

et al., 2010). For the less water-soluble compounds, increased emissions reduced the bioavailability, and increases in temperature resulted in an overall reduction in net bioaccumulation (Borga et al., 2010).

8.2. Long-term trends

Biota data in the polar regions were collected from literature that covered different time periods and different statistical analysis methods to build long-term trends (Cincinelli et al., 2016; Ellis et al., 2018; Helgason et al., 2008; Riget et al., 2010a; Riget et al., 2019). The longest time span involved the reporting of POPs in penguins for nearly 50 years (1964–2011) in the Antarctic (Ellis et al., 2018), but other long-term studies (over 20–30 years) have also been conducted (Cincinelli et al., 2016; Goerke et al., 2004; Helgason et al., 2008; Riget et al., 2010a). The biota chosen to reflect the long-term trends of POPs mainly include eggs, fat tissues in penguins, fish, and *Trematomus bernacchii* (AMAP, 2016; Braune, 2007; Ellis et al., 2018; Goerke et al., 2004; Helgason et al., 2008; Riget et al., 2010a; Riget et al., 2010b, 2019). Generally, legacy POP levels in most species increased from the 1960s–1980s, and then decreased from the 1990s to the present day, with only very few cases of significantly increasing trends having been reported (Braune and Letcher, 2013; Cincinelli et al., 2016; Ellis et al., 2018; Goerke et al., 2004; Helgason et al., 2008; Riget et al., 2010a).

The temporal trends of POPs in biota are influenced by the expression of their concentrations, of food webs (pelagic or benthic), and of species and locations (Evenset et al., 2016; Goutte et al., 2013; Lescord et al., 2015; Moisey et al., 2001; Riget et al., 2010b). For example, Riget et al. (2010b) found decreasing trends of POPs during the period 1994–2008 in Arctic Char became non-significant when expressed by lipid weight rather than wet weight. Significantly increasing concentrations within a decade (1987–1996) were observed for two benthic fish species, while the pelagic species did not increase (Goerke et al., 2004). van den Brink et al. (2011) concluded that concentrations of PCBs, DDE, HCB and dieldrin were declining during 1993–2003 in Antarctic pelagic seabirds, but not in benthic organisms. Braune (2007) measured OCPs and PCBs in eggs of three species (murrelets, fulmars and kittiwakes) in the Arctic, with the temporal trends for HCH (1975–2003) found to be different among the species. Riget et al. (2010a) observed that the highest mean decreasing rate in Arctic biota was for α -HCH, γ -HCH and DDTs, while the lowest rate was for total chlordanes and toxaphene. Moreover, β -HCH showed a low or nonexistent rate of decline in the western Arctic, but greater decreasing trends to the east (Riget et al., 2010a).

8.3. Exposure assessment

The bioconcentration and toxicity of POPs result in high exposure risks for species at the top trophic levels (Allen-Gil et al., 1997; Andersen et al., 2001; Braune et al., 2005; Bustnes et al., 2007; Goerke et al., 2004; Hoekstra et al., 2003; Hop et al., 2002; Kelly and Gobas, 2001; Kelly et al., 2008; Rudolph et al., 2016; Villa et al., 2017). Moreover, the natural stresses on biota in the extreme environments of the polar regions are often more severe than in other regions (<https://www.amap.no/documents/doc/amap-assessment-2018-biological-effects-of-contaminants-on-arctic-wildlife-and-fish/1663>, AMAP, 2018), rendering species here more vulnerable to the effects of pollutants.

Comparatively, the Arctic has attracted more attention, due to the traditional lifestyle of the Inuit population appearing to be an important contributor to POP exposure (Dallaire et al., 2009; Valera et al., 2013; Zhulidov et al., 2002). Letcher et al. (2010) and <https://www.amap.no/documents/doc/amap-assessment-2015-human->

health-in-the-arctic/1346, AMAP (2015) reviewed the biological effects in relation to organohalogen contaminant exposure in several top trophic level species of the Arctic, and tried to assess known tissue/body compartment concentration data in the context of possible threshold levels of effects to evaluate the risks. Dallaire et al. (2009) found the consumption of fish and marine mammals appears to be an important contributor to PFOS and PBDE exposure among Nunavik Inuit. Valera et al. (2013) concluded that some PCB congeners and p,p'-DDE are associated with a higher risk of hypertension in Inuit adults.

Only small amounts of exposure-assessment research in the Antarctic and TP have been reported (Borghesi et al., 2008; Wang et al., 2016b). In the Antarctic, the TEQ values of PCBs and OCPs, and the “toxic equivalent factors” of PCDF congeners, have been used to estimate exposure risks (Borghesi et al., 2008; Schiavone et al., 2009). The TEQ values were found to be one order of magnitude lower than those considered to elicit physiological effects in aquatic mammals (Borghesi et al., 2008; Schiavone et al., 2009). Generally, POP concentrations are not currently a risk in terms of generating adverse health effects. In the TP, the exposure risks with respect to POPs have mainly been examined in the context of human beings Bi et al. (2016); Chen et al. (2015); Gong et al. (2011); Liu et al. (2014); Pan et al. (2014); Pokhrel et al. (2018); Wang et al. (2016a); Wang et al. (2016b). The results showed that OCPs in farmland areas do not pose any non-carcinogenic risks, and only pose a very low cancer risk to Tibetan residents (Wang et al., 2016b).

9. Modeling

The presence of POPs in the polar regions derive primarily from external inputs, as evidenced by the numerous studies mentioned above, for all environmental media (atmosphere, water, soil, sediment, and biota). However, although the transport pathways of POPs from source regions to the polar regions have been observed, the specific mechanisms of transportation and the environmental burdens of these substances are harder to determine based on observational data alone. Thus, models involving multi-compartmental cycling are needed to simulate the transport mechanisms and verify the fate of POPs in the polar regions. Several models have been developed to examine the transport and transfer of POPs from low latitudes to the Arctic, but very few have been developed to investigate the equivalent the Antarctic and TP.

Based on the direction of transport (i.e., horizontal and vertical), both 2-D and 3-D transport models have been developed (Christensen, 1993; Krapivin, 1995; Wania and Mackay, 1995). Models concerning the transport of POPs to the Arctic are as follows: (1) global atmospheric distribution models for POPs (Wania and Dugani, 2003; Wania and Mackay, 1993; Wania and Mackay, 1995; Wania and Mackay, 1999; Wania et al., 1999); (2) the Danish Eulerian Hemispheric Model (Friedman and Selin, 2016; Hansen et al., 2004); (3) the Arctic Mass Balance Box Model (Li et al., 2004); GEM/POP (Huang et al., 2007); (4) ocean transport model (Armitage et al., 2006; Armitage et al., 2009; Stemmler and Lammel, 2009; Stemmler and Lammel, 2010; Stemmler and Lammel, 2013; Zhang et al., 2017); (5) a coupled atmosphere–ocean general circulation model (Guglielmo et al., 2009); (6) a 2-D atmospheric transport model (Ma, 2010); (7) a 3-D atmospheric transport model (Zhang et al., 2010); and (8) GEOS-Chem (Friedman and Selin, 2012; Friedman and Selin, 2016).

Some atmospheric transport models have been used to estimate the transmission height and distance for POPs from source regions (Asia, Europe and North Atlantic) to the Arctic (Halsall et al., 2001; Ma, 2010; Zhang et al., 2010; Octaviani et al., 2015). For example, the poleward atmospheric transports of γ -HCH from Asian or

Eurasian sources to the Arctic have been found to occur mostly in the mid-troposphere (e.g., from 3000 m to 5500 m) (Zhang et al., 2010). Huang et al. (2007) used GEM/POPs to demonstrate that, for PCBs, transport flux peaking occurs below 8 km for the gaseous phase and 14 (6) km for particulate PCB-28 (PCB-180).

The transport mass of POPs in the Arctic has also been estimated by models. For instance, Huang et al. (2007) estimated the Arctic receives 5914 kg (89 kg) of PCB-28 (PCB-180) from Europe, 1217 kg (26 kg) from the North Atlantic, and exports 2413 kg (8.7 kg) of PCB-28 (PCB-180) to North America, and 21.4 kg PCB-180 to Asia in year 2000. Between 1945 and 2000, the total loading of α -HCH to the Arctic Ocean was approximately 27 700 t, accounting for approximately 0.6% of all global α -HCH emissions from agricultural land to the atmosphere (Li et al., 2004). Similar results were reported by Wania and Mackay (1999), who found only very small fractions of α -HCH (<1%) have reached the Arctic. The dominant input pathways for α -HCH entering the Arctic are gas absorption (50%) and ocean currents (34%); and the major means of removal of α -HCH are microbial degradation (49%) and ocean currents (23%) (Li et al., 2004).

Climate change also impacts the atmospheric transport and fate of POPs in the Arctic (Hansen et al., 2015; Lamon et al., 2009; Ma et al., 2016; Ma et al., 2011). By the end of the 2090s, the mass of HCHs within the Arctic is predicted to be up to 38% higher than that in the 1990s, whereas the change in the mass of PCBs is expected to range from 38% lower to 17% higher (Hansen et al., 2015). Emerging contaminants, i.e., those with no or only a short emissions history, will be more rapidly transported to, and build up within, the Arctic environment in the future warmer climate (Hansen et al., 2015). Considering the influence of climate- and emissions-driven processes on Arctic PCB concentrations, Friedman and Selin (2016) evidenced that the emission or removal/transformation processes occurring outside the Arctic, rather than within, have a greater influence on atmospheric PCB levels.

10. Research needs and future perspectives

Although the levels of atmospheric POPs have been reported for the polar regions, the temporal trends and half-lives for specific compounds, and the role that atmospheric circulation plays in the transport of “emerging” POPs, remain unclear for the Antarctic and the TP. Thus, long-term and consistent monitoring of atmospheric POPs is required in the polar regions.

The monitoring of POPs in water has yielded data for more than 20 years in the Arctic and Antarctic, whereas related work in the TP has only just begun. The TP is also often referred to as the “Asia water tower”, because it is home to the most mountain glaciers in the world, as well as thousands of lakes on the Tibetan Plateau, both of which are vital to Asia water security and beyond. In this context, more observations of POPs in water in the TP are needed to evaluate the global cycling of POPs.

Systematic research has been performed on the levels and bio-accumulation of POPs in the aquatic food webs of the polar regions. Comparatively, however, fewer data have been published on terrestrial food chains, both in terms of the quantity and length of the chains, because only two short terrestrial food chains (lichen–caribou–wolf in the Arctic and air–grass–yak in the TP) have thus far been reported. Due to close association that terrestrial food chains have with human life, more work on the bioaccumulation mechanism from the environment to higher trophic levels is urgently needed in the future.

So far, results have indicated that POPs in the Antarctic and TP do not cause adverse effects on animals or humans; whereas, in the Arctic, the traditional lifestyle of the Inuit population has not protected them against exposure to POPs, particularly emerging

pollutants. Continuous attention should be paid to the health risk of POPs on the Inuit and Tibetan people. Regardless, there remains a dearth of data on the true nature of contaminant exposure, especially the cause–effect relationships with respect to the exposure of these contaminants in wildlife and people.

Models have been used to simulate the global transport of POPs to the Arctic, but models that simulate the transport processes and masses of POPs in the Antarctic and TP should be developed. Moreover, it will be challenging to develop models that cover the complexity of the transport processes (including glacial melt) for POPs across the Himalaya and in regions that experience strong solar radiation.

Climate change and emissions reduction measures act jointly on the fate of POPs. In this regard, the polar regions are highly sensitive to climatic factors but far away from direct emission sources, thus offering a natural laboratory to carry out research on the effects of both climate and emissions on the fate of POPs. However, presently, studies are limited by a lack of long-term and consistent data, making it difficult to fully understand the mechanisms involved, especially in the Antarctic and TP. Hence, the long-term monitoring of POPs together with the development of emissions inventories are a necessary next step.

Finally, although the environmental processes related to POPs have been separately observed in each polar region, there remains no consensus on the fate of POPs in remote and cold regions. Associated work in the three polar regions on all environmental media (air, soil, water and biota) should be placed on the agenda.

Acknowledgements

This study was supported by the Strategic Priority Research Program of Chinese Academy of Sciences, Pan-Third Pole Environment Study for a Green Silk Road (Pan-TPE) (XDA2004050202), the National Natural Science Foundation of China (41671480 and 41701083), and China Postdoctoral Science Foundation (2018T110146).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.envpol.2019.01.093>.

References

- Aas, C.B., Fuglei, E., Herzke, D., Yoccoz, N.G., Routti, H., 2014. Effect of body condition on tissue distribution of perfluoroalkyl substances (PFASs) in arctic fox (*Vulpes lagopus*). *Environ. Sci. Technol.* 48, 11654–11661.
- Abakumov, E.V., Lodygin, E.D., Gabov, D.A., Krylenkov, V.A., 2014. Polycyclic aromatic hydrocarbons content in Antarctica soils as exemplified by the Russian polar stations. *Gigiena i sanitarii* 31–35.
- Ahrens, L., Gerwinski, W., Theobald, N., Ebinghaus, R., 2010. Sources of polyfluoroalkyl compounds in the North Sea, Baltic Sea and Norwegian Sea: evidence from their spatial distribution in surface water. *Mar. Pollut. Bull.* 60, 255–260.
- Aislabe, J., Balks, M., Astori, N., Stevenson, G., Symons, R., 1999. Polycyclic aromatic hydrocarbons in fuel-oil contaminated soils, Antarctica. *Chemosphere* 39, 2201–2207.
- Alexeeva, L.B., Strachan, W.M.J., Shlychkova, V.V., Nazarova, A.A., Nikanorov, A.M., Korotova, L.G., Koreneva, V.I., 2001. Organochlorine pesticide and trace metal monitoring of Russian rivers flowing to the Arctic Ocean: 1990–1996. *Mar. Pollut. Bull.* 43, 71–85.
- Allen-Gil, S.M., Gubala, C.P., Wilson, R., Landers, D.H., Wade, T.L., Sericano, J.L., Curtis, L.R., 1997. Organochlorine pesticides and polychlorinated biphenyls (PCBs) in sediments and biota from four US Arctic lakes. *Arch. Environ. Contam. Toxicol.* 33, 378–387.
- AMAP, 1998. AMAP Assessment 1998. Chapter 6. Persistent Organic Pollutants. Arctic Monitoring and Assessment Programme (AMAP), Oslo, Norway, pp. 183–335.
- AMAP, 2004. AMAP Assessment 2002: Persistent Organic Pollutants in the Arctic. Arctic Monitoring and Assessment Programme (AMAP), Oslo, Norway xvi + 310.
- AMAP, 2009. AMAP Assessment 2009: Human Health in the Arctic. Arctic Monitoring and Assessment Programme, Oslo, Norway xiv + 254.
- AMAP, 2015. AMAP Assessment 2015: Human Health in the Arctic. Arctic Monitoring and Assessment Programme, Oslo, Norway xii + 165.
- AMAP, 2016. AMAP Assessment 2015: Temporal Trends in Persistent Organic Pollutants in the Arctic. Arctic Monitoring and Assessment Programme, Oslo, Norway vi + 71.
- AMAP, 2017. AMAP Assessment 2016: Chemicals of Emerging Arctic Concern. Arctic Monitoring and Assessment Programme (AMAP), Oslo, Norway xvi + 353.
- AMAP, 2018. AMAP Assessment 2018: Biological Effects of Contaminants on Arctic Wildlife and Fish. Arctic Monitoring and Assessment Programme (AMAP), Oslo, Norway vii + 84.
- Andersen, G., Kovacs, K.M., Lydersen, C., Skaare, J.U., Gjert, I., Jenssen, B.M., 2001. Concentrations and patterns of organochlorine contaminants in white whales (*Delphinapterus leucas*) from Svalbard, Norway. *Sci. Total Environ.* 264, 267–281.
- Andersson, P.L., Berg, A.H., Olsson, P.E., Tysklind, M., 1998. Distribution of selected polychlorinated biphenyls (PCBs) in brain and liver of Arctic char (*Salvelinus alpinus*). *Mar. Environ. Res.* 46, 501–504.
- Armitage, J., Cousins, I.T., Buck, R.C., Prevedouros, K., Russell, M.H., MacLeod, M., Korzeniowski, S.H., 2006. Modeling global-scale fate and transport of perfluorooctanoate emitted from direct sources. *Environ. Sci. Technol.* 40, 6969–6975.
- Armitage, J.M., MacLeod, M., Cousins, I.T., 2009. Modeling the global fate and transport of perfluorooctanoic acid (PFOA) and perfluorooctanoate (PFO) emitted from direct sources using a multispecies mass balance model. vol 43, pg 1134, 2009 *Environ. Sci. Technol.* 43, 6438–6439.
- Augusto, S., Máguas, C., Branquinho, C., 2013. Guidelines for biomonitoring persistent organic pollutants (POPs), using lichens and aquatic mosses – a review. *Environ. Pollut.* 180, 330–338.
- Barber, J.L., Thomas, G.O., Kerstiens, G., Jones, K.C., 2004. Current issues and uncertainties in the measurement and modelling of air-vegetation exchange and within-plant processing of POPs. *Environ. Pollut.* 128, 99–138.
- Baek, S.-Y., Choi, S.-D., Chang, Y.-S., 2011. Three-year atmospheric monitoring of organochlorine pesticides and polychlorinated biphenyls in polar regions and the south pacific. *Environ. Sci. Technol.* 45, 4475–4482.
- Becker, L., Scheringer, M., Schenker, U., Hungerbuehler, K., 2011. Assessment of the environmental persistence and long-range transport of endosulfan. *Environ. Pollut.* 159, 1737–1743.
- Benskin, J.P., Ahrens, L., Muir, D.C.G., Scott, B.F., Spencer, C., Rosenberg, B., Tomy, G., Kylin, H., Lohmann, R., Martin, J.W., 2012a. Manufacturing origin of perfluorooctanoate (PFOA) in Atlantic and Canadian Arctic seawater. *Environ. Sci. Technol.* 46, 677–685.
- Benskin, J.P., Muir, D.C.G., Scott, B.F., Spencer, C., De Silva, A.O., Kylin, H., Martin, J.W., Morris, A., Lohmann, R., Tomy, G., Rosenberg, B., Taniyasu, S., Yamashita, N., 2012b. Perfluoroalkyl acids in the Atlantic and Canadian Arctic oceans. *Environ. Sci. Technol.* 46, 5815–5823.
- Bi, X., Luo, W., Gao, J., Xu, L., Guo, J., Zhang, Q., Romesh, K.Y., Giesy, J.P., Kang, S., de Boer, J., 2016. Polycyclic aromatic hydrocarbons in soils from the Central-Himalaya region: distribution, sources, and risks to humans and wildlife. *Sci. Total Environ.* 556, 12–22.
- Bidleman, T.F., 1999. Atmospheric transport and air-surface exchange of pesticides. *Water Air Soil Pollut.* 115, 115–166.
- Bidleman, T.F., Kylin, H., Jantunen, L.M., Helm, P.A., MacDonald, R.W., 2007. Hexachlorocyclohexanes in the Canadian Archipelago. 1. Spatial distribution and pathways of alpha-, beta- and gamma-HCHs in surface water. *Environ. Sci. Technol.* 41, 2688–2695.
- Bigot, M., Hawker, D.W., Cropp, R., Muir, D.C.G., Jensen, B., Bossi, R., Nash, S.M.B., 2017. Spring melt and the redistribution of organochlorine pesticides in the sea-ice environment: a comparative study between Arctic and Antarctic regions. *Environ. Sci. Technol.* 51, 8944–8952.
- Bigot, M., Muir, D.C.G., Hawker, D.W., Cropp, R., Dachs, J., Teixeira, C.F., Nash, S.B., 2016. Air-seawater exchange of organochlorine pesticides in the Southern Ocean between Australia and Antarctica. *Environ. Sci. Technol.* 50, 8001–8009.
- Bizzotto, E.C., Villa, S., Vaj, C., Vighi, M., 2009. Comparison of glacial and non-glacial-fed streams to evaluate the loading of persistent organic pollutants through seasonal snow/ice melt. *Chemosphere* 74, 924–930.
- Blais, J.M., Schindler, D.W., Muir, D.C.G., Kimpe, L.E., Donald, D.B., Rosenberg, B., 1998. Accumulation of persistent organochlorine compounds in mountains of western Canada. *Nature* 395, 585–588.
- Blais, J.M., Schindler, D.W., Muir, D.C.G., Sharp, M., Donald, D., Lafreniere, M., Braekvelt, E., Strachan, W.M.J., 2001. Melting glaciers: a major source of persistent organochlorines to subalpine Bow Lake in Banff National Park, Canada. *Ambio* 30, 410–415.
- Borga, K., Saloranta, T.M., Ruus, A., 2010. Simulating climate change-induced alterations in bioaccumulation of organic contaminants in an Arctic marine food web. *Environ. Toxicol. Chem.* 29, 1349–1357.
- Borghesi, N., Corsolini, S., Focardi, S., 2008. Levels of polybrominated diphenyl ethers (PBDEs) and organochlorine pollutants in two species of Antarctic fish (*Chionodraco hamatus* and *Trematomus bernacchii*). *Chemosphere* 73, 155–160.
- Borghini, F., Grimalt, J.O., Sanchez-Hernandez, J.C., Bargagli, R., 2005a. Organochlorine pollutants in soils and mosses from Victoria Land (Antarctica). *Chemosphere* 58, 271–278.
- Borghini, F., Grimalt, J.O., Sanchez-Hernandez, J.C., Barra, R., Garcia, C.J.T., Focardi, S., 2005b. Organochlorine compounds in soils and sediments of the mountain

- Andean Lakes. *Environ. Pollut.* 136, 253–266.
- Bossi, R., Skjoth, C.A., Skov, H., 2013. Three years (2008–2010) of measurements of atmospheric concentrations of organochlorine pesticides (OCPs) at Station Nord, North-East Greenland. *Environmental Science-Processes & Impacts* 15, 2213–2219.
- Braune, B., Muir, D., DeMarch, B., Gamberg, M., Poole, K., Currie, R., Dodd, M., Dushenko, W., Eamer, J., Elkin, B., Evans, M., Grundy, S., Hebert, C., Johnstone, R., Kidd, K., Koenig, B., Lockhart, L., Marshall, H., Reimer, K., Sanderson, J., Shutt, L., 1999. Spatial and temporal trends of contaminants in Canadian Arctic freshwater and terrestrial ecosystems: a review. *Sci. Total Environ.* 230, 145–207.
- Braune, B.A., 2007. Temporal trends of organochlorines and mercury in seabird eggs from the Canadian Arctic, 1975–2003. *Environ. Pollut.* 148, 599–613.
- Braune, B.M., Donaldson, G.M., Hobson, K.A., 2002. Contaminant residues in seabird eggs from the Canadian Arctic. II. Spatial trends and evidence from stable isotopes for intercolony differences. *Environ. Pollut.* 117, 133–145.
- Braune, B.M., Letcher, R.J., 2013. Perfluorinated sulfonate and carboxylate compounds in eggs of seabirds breeding in the Canadian arctic: temporal trends (1975–2011) and interspecies comparison. *Environ. Sci. Technol.* 47, 616–624.
- Braune, B.M., Letcher, R.J., Gaston, A.J., Mallory, M.L., 2015. Trends of polybrominated diphenyl ethers and hexabromocyclodecane in eggs of Canadian Arctic seabirds reflect changing use patterns. *Environ. Res.* 142, 651–661.
- Braune, B.M., Mallory, M.L., 2017. Declining trends of polychlorinated dibenzo-p-dioxins, dibenzofurans and non-ortho PCBs in Canadian Arctic seabirds. *Environ. Pollut.* 220, 557–566.
- Braune, B.M., Outridge, P.M., Fisk, A.T., Muir, D.C.G., Helm, P.A., Hobbs, K., Hoekstra, P.F., Kozuk, Z.A., Kwan, M., Letcher, R.J., Lockhart, W.L., Norstrom, R.J., Stern, G.A., Stirling, I., 2005. Persistent organic pollutants and mercury in marine biota of the Canadian Arctic: an overview of spatial and temporal trends. *Sci. Total Environ.* 351, 4–56.
- Bright, D.A., Dushenko, W.T., Grundy, S.L., Reimer, K.J., 1995. Evidence for short-range transport of polychlorinated-biphenyls in the Canadian arctic using signatures of PCBs in soils. *Sci. Total Environ.* 160–61, 251–263.
- Burkow, I.C., Kallenborn, R., 2000. Sources and transport of persistent pollutants to the Arctic. *Toxicol. Lett.* 112, 87–92.
- Busch, J., Ahrens, L., Xie, Z., Sturm, R., Ebinghaus, R., 2010. Polyfluoroalkyl compounds in the East Greenland Arctic Ocean. *J. Environ. Monit.* 12, 1242–1246.
- Bustnes, J.O., Tveraa, T., Varpe, O., Henden, J.A., Skaare, J.U., 2007. Reproductive performance and organochlorine pollutants in an Antarctic marine top predator: the south polar skua. *Environ. Int.* 33, 911–918.
- Cabrero, A., Dachs, J., Barcelo, D., Jones, K.C., 2013. Climatic and biogeochemical controls on the remobilization and reservoirs of persistent organic pollutants in Antarctica. *Environ. Sci. Technol.* 47, 4299–4306.
- Cabrero, A., Dachs, J., Barcelo, D., Jones, K.C., 2012. Influence of organic matter content and human activities on the occurrence of organic pollutants in antarctic soils, lichens, grass, and mosses. *Environ. Sci. Technol.* 46, 1396–1405.
- Cabrero, A., Larramendi, R., Albar, J.-P., Dachs, J., 2017. Persistent organic pollutants in the atmosphere of the Antarctic Plateau. *Atmos. Environ.* 149, 104–108.
- Cai, M., Ma, Y., Xie, Z., Zhong, G., Moeller, A., Yang, H., Sturm, R., He, J., Ebinghaus, R., Meng, X.-Z., 2012a. Distribution and air-sea exchange of organochlorine pesticides in the North Pacific and the Arctic. *Journal of Geophysical Research-Atmospheres* 117.
- Cai, M., Qiu, C., Shen, Y., Cai, M., Huang, S., Qian, B., Sun, J., Liu, X., 2010. Concentration and distribution of 17 organochlorine pesticides (OCPs) in seawater from the Japan Sea northward to the Arctic Ocean. *Sci. China Chem.* 53, 1033–1047.
- Cai, M., Zhao, Z., Yin, Z., Ahrens, L., Huang, P., Cai, M., Yang, H., He, J., Sturm, R., Ebinghaus, R., Xie, Z., 2012b. Occurrence of perfluoroalkyl compounds in surface waters from the North Pacific to the Arctic ocean. *Environ. Sci. Technol.* 46, 661–668.
- Cai, M.G., Hong, Q.Q., Wang, Y., Luo, X.J., Chen, S.J., Cai, M.H., Qiu, C.R., Huang, S.Y., Mai, B.X., 2012c. Distribution of polybrominated diphenyl ethers and decabromodiphenylethane in surface sediments from the Bering Sea, Chukchi Sea, and Canada Basin. *Deep-Sea Res. Part II Top. Stud. Oceanogr.* 81–84, 95–101.
- Cannon, S.M.E., Lavers, J.L., Figueiredo, B., 2016. Plastic ingestion by fish in the Southern Hemisphere: a baseline study and review of methods. *Mar. Pollut. Bull.* 107, 286–291.
- Carlsson, P., Cornelissen, G., Boggild, C.E., Rysgaard, S., Mortensen, J., Kallenborn, R., 2012. Hydrology-linked spatial distribution of pesticides in a fjord system in Greenland. *J. Environ. Monit.* 14, 1437–1443.
- Carrizo, D., Gustafsson, O., 2011a. Distribution and inventories of polychlorinated biphenyls in the polar mixed layer of seven pan-arctic shelf seas and the interior basins. *Environ. Sci. Technol.* 45, 1420–1427.
- Carrizo, D., Sobek, A., Salvado, J.A., Gustafsson, O., 2017. Spatial distributions of DDTs in the water masses of the Arctic ocean. *Environ. Sci. Technol.* 51, 7913–7919.
- Carroll, J., Savinov, V., Savinova, T., Dahle, S., McCrea, R., Muir, D.C.G., 2008. PCBs, PBDEs and pesticides released to the Arctic ocean by the Russian rivers Ob and Yenisei. *Environ. Sci. Technol.* 42, 69–74.
- Carrizo, D., Gustafsson, O., 2011b. Pan-arctic river fluxes of polychlorinated biphenyls. *Environ. Sci. Technol.* 45, 8377–8384.
- Casal, P., Castro-Jiménez, J., Pizarro, M., Katsoyiannis, A., Dachs, J., 2018. Seasonal soil/snow-air exchange of semivolatile organic pollutants at a coastal arctic site (Tromsø, 69°N). *Sci. Total Environ.* 636, 1109–1116.
- Chakraborty, P., Khuman, S.N., Selvaraj, S., Sampath, S., Devi, N.L., Bang, J.J., Katsoyiannis, A., 2016. Polychlorinated biphenyls and organochlorine pesticides in river brahmaputra from the outer himalayan range and river hooghly emptying into the bay of bengal: occurrence, sources and ecotoxicological risk assessment. *Environ. Pollut.* 219, 998–1006.
- Chen, D.Z., Liu, W.J., Liu, X.D., Westgate, J.N., Wania, F., 2008. Cold-Trapping of persistent organic pollutants in the mountain soils of western sichuan, China. *Environ. Sci. Technol.* 42, 9086–9091.
- Chen, P.F., Kang, S.C., Li, C.L., Rupakheti, M., Yan, F.P., Li, Q.L., Ji, Z.M., Zhang, Q.G., Luo, W., Sillanpää, M., 2015. Characteristics and sources of polycyclic aromatic hydrocarbons in atmospheric aerosols in the Kathmandu Valley, Nepal. *Sci. Total Environ.* 538, 86–92.
- Cheng, H., Lin, T., Zhang, G., Liu, G., Zhang, W., Qi, S., Jones, K.C., Zhang, X., 2014a. DDTs and HCHs in sediment cores from the Tibetan Plateau. *Chemosphere* 94, 183–189.
- Cheng, H.R., Lin, T., Zhang, G., Liu, G.Q., Zhang, W.L., Qi, S.H., Jones, K.C., Zhang, X.W., 2014b. DDTs and HCHs in sediment cores from the Tibetan Plateau. *Chemosphere* 94, 183–189.
- Choi, S.-D., Baek, S.-Y., Chang, Y.S., Wania, F., Ikononou, M.G., Yoon, Y.-J., Park, B.-K., Hong, S., 2008. Passive air sampling of polychlorinated biphenyls and organochlorine pesticides at the Korean Arctic and Antarctic research stations: implications for long-range transport and local pollution. *Environ. Sci. Technol.* 42, 7125–7131.
- Christensen, J., 1993. Testing advection schemes in a 3-dimensional air-pollution model. *Math. Comput. Model.* 18, 75–88.
- Cincinelli, A., Martellini, T., Del Bubba, M., Lepri, L., Corsolini, S., Borghesi, N., King, M.D., Dickhut, R.M., 2009. Organochlorine pesticide air-water exchange and bioconcentration in krill in the Ross Sea. *Environ. Pollut.* 157, 2153–2158.
- Cincinelli, A., Martellini, T., Pozo, K., Kukucka, P., Audy, O., Corsolini, S., 2016. Trematomus bernacchii as an indicator of POP temporal trend in the Antarctic seawaters. *Environ. Pollut.* 217, 19–25.
- Cipro, C.V.Z., Taniguchi, S., Montone, R.C., 2017. Organic pollutants in snow and seasonal melting water from king George island, Antarctica. *Water Air Soil Pollut.* 228.
- Combi, T., Martins, C.C., Taniguchi, S., Leonel, J., Lourenco, R.A., Montone, R.C., 2017. Depositional history and inventories of polychlorinated biphenyls (PCBs) in sediment cores from an antarctic specially managed area (admiralty bay, king George island). *Mar. Pollut. Bull.* 118, 447–451.
- Corsolini, S., 2009. Industrial contaminants in Antarctic biota. *J. Chromatogr. A* 1216, 598–612.
- Corsolini, S., Ademollo, N., Martellini, T., Randazzo, D., Vacchi, M., Cincinelli, A., 2017. Legacy persistent organic pollutants including PBDEs in the trophic web of the Ross Sea (Antarctica). *Chemosphere* 185, 699–708.
- Corsolini, S., Ademollo, N., Romeo, T., Olmastroni, S., Focardi, S., 2003. Persistent organic pollutants in some species of a Ross Sea pelagic trophic web. *Antarct. Sci.* 15, 95–104.
- Corsolini, S., Kannan, K., Imagawa, T., Focardi, S., Giesy, J.P., 2002. Polychloronaphthalenes and other dioxin-like compounds in Arctic and Antarctic marine food webs. *Environ. Sci. Technol.* 36, 3490–3496.
- Corsolini, S., Sara, G., 2017. The trophic transfer of persistent pollutants (HCB, DDTs, PCBs) within polar marine food webs. *Chemosphere* 177, 189–199.
- Court, G.S., Davies, L.S., Focardi, S., Bargagli, R., Fossi, C., Leonzio, C., Marili, L., 1997. Chlorinated hydrocarbons in the tissues of south polar skuas (*Catharacta maccormicki*) and adelia penguins (*pygoscelis adeliae*) from ross sea, Antarctica. *Environ. Pollut.* 97, 295–301.
- Dachs, J., Lohmann, R., Ockenden, W.A., Mejanelle, L., Eisenreich, S.J., Jones, K.C., 2002. Oceanic biogeochemical controls on global dynamics of persistent organic pollutants. *Environ. Sci. Technol.* 36, 4229–4237.
- Dahle, S., Savinov, V.M., Matishov, G.G., Evensen, A., Naes, K., 2003. Polycyclic aromatic hydrocarbons (PAHs) in bottom sediments of the Kara sea shelf, Gulf of Ob and Yenisei bay. *Sci. Total Environ.* 306, 57–71.
- Dallaire, R., Ayotte, P., Pereg, D., Dery, S., Dumas, P., Langlois, E., Dewailly, E., 2009. Determinants of plasma concentrations of perfluorooctanesulfonate and brominated organic compounds in Nunavut Inuit adults (Canada). *Environ. Sci. Technol.* 43, 5130–5136.
- Daly, G.L., Lei, Y.D., Teixeira, C., Muir, D.C.G., Wania, F., 2007. Pesticides in western Canadian mountain air and soil. *Environ. Sci. Technol.* 41, 6020–6025.
- De Laender, F., Van Oevelen, D., Frantzen, S., Middelburg, J.J., Soetaert, K., 2010. Seasonal PCB bioaccumulation in an arctic marine ecosystem: a model analysis incorporating lipid dynamics, food-web productivity and migration. *Environ. Sci. Technol.* 44, 356–361.
- Dietz, R., Riget, F., Cleemann, M., Aarkrog, A., Johansen, P., Hansen, J.C., 2000. Comparison of contaminants from different trophic levels and ecosystems. *Sci. Total Environ.* 245, 221–231.
- Ding, X., Wang, X.-M., Xie, Z.-Q., Xiang, C.-H., Mai, B.-X., Sun, L.-G., Zheng, M., Sheng, G.-Y., Fu, J.-M., 2007. Atmospheric hexachlorocyclohexanes in the North Pacific Ocean and the adjacent Arctic region: spatial patterns, chiral signatures, and sea-air exchanges. *Environ. Sci. Technol.* 41, 5204–5209.
- Dong, C., Bai, X., Sheng, H., Jiao, L., Zhou, H., Shao, Z., 2015. Distribution of PAHs and the PAH-degrading bacteria in the deep-sea sediments of the high-latitude Arctic Ocean. *Biogeosciences* 12, 2163–2177.
- Duan, Y., Wu, B.X., Xu, L., He, J.X., Sun, T., 2011. Characterisation of n-alkanes and their hydrogen isotopic composition in sediments from Lake Qinghai, China. *Org. Geochem.* 42, 720–726.
- Ellis, D.S., Cipro, C.V.Z., Ogletree, C.A., Smith, K.E., Aronson, R.B., 2018. A 50-year retrospective of persistent organic pollutants in the fat and eggs of penguins of the Southern Ocean. *Environ. Pollut.* 241, 155–163.

- Evenset, A., Hallanger, I.G., Tessmann, M., Warner, N., Ruus, A., Borga, K., Gabrielsen, G.W., Christensen, G., Renaud, P.E., 2016. Seasonal variation in accumulation of persistent organic pollutants in an Arctic marine benthic food web. *Sci. Total Environ.* 542, 108–120.
- Fellin, P., Barrie, L.A., Dougherty, D., Toom, D., Muir, D., Grift, N., Lockhart, L., Billeck, B., 1996. Air monitoring in the arctic: results for selected persistent organic pollutants for 1992. *Environ. Toxicol. Chem.* 15, 253–261.
- Fernandez, P., Carrera, G., Grimalt, J.O., 2005. Persistent organic pollutants in remote freshwater ecosystems. *Aquat. Sci.* 67, 263–273.
- Fernandez, P., Vilanova, R.M., Grimalt, J.O., 1999. Sediment fluxes of polycyclic aromatic hydrocarbons in European high altitude mountain lakes. *Environ. Sci. Technol.* 33, 3716–3722.
- Fernandez, P., Vilanova, R.M., Martinez, C., Appleby, P., Grimalt, J.O., 2000. The historical record of atmospheric pyrolytic pollution over Europe registered in the sedimentary PAH from remote mountain lakes. *Environ. Sci. Technol.* 34, 1906–1913.
- Fisk, A.T., Hobson, K.A., Norstrom, R.J., 2001. Influence of chemical and biological factors on trophic transfer of persistent organic pollutants in the northwestern polynya marine food web. *Environ. Sci. Technol.* 35, 732–738.
- Franzaring, J., van der Eerden, L.J.M., 2000. Accumulation of airborne persistent organic pollutants (POPs) in plants. *Basic Appl. Ecol.* 1, 25–30.
- Friedman, C.L., Selin, N.E., 2012. Long-range atmospheric transport of polycyclic aromatic hydrocarbons: a global 3-D model analysis including evaluation of arctic sources. *Environ. Sci. Technol.* 46, 9501–9510.
- Friedman, C.L., Selin, N.E., 2016. PCBs in the Arctic atmosphere: determining important driving forces using a global atmospheric transport model. *Atmos. Chem. Phys.* 16, 3433–3448.
- Fromberg, A., Cleemann, M., Carlsen, L., 1999. Review on persistent organic pollutants in the environment of Greenland and Faroe Islands. *Chemosphere* 38, 3075–3093.
- Fu, S., Chu, S., Xu, X., 2001. Organochlorine pesticide residue in soils from Tibet, China. *Bull. Environ. Contam. Toxicol.* 66, 171–177.
- Fuoco, R., Giannarelli, S., Onor, M., Ghimenti, S., Abete, C., Termine, M., Francesconi, S., 2012. A snow/firm four-century record of polycyclic aromatic hydrocarbons (PAHs) and polychlorobiphenyls (PCBs) at Talos Dome (Antarctica): *Microchem. J.* 105, 133–141.
- Galassi, S., Valsecchi, S., Tartari, G.A., 1997. The distribution of PCB's and chlorinated pesticides in two connected Himalayan lakes. *Water Air Soil Pollut.* 99, 717–725.
- Galban-Malagon, C., Berrojalbiz, N., Ojeda, M.J., Dachs, J., 2012. The oceanic biological pump modulates the atmospheric transport of persistent organic pollutants to the Arctic. *Nat. Commun.* 3.
- Galban-Malagon, C.J., Berrojalbiz, N., Gioia, R., Dachs, J., 2013. The "degradative" and "biological" pumps controls on the atmospheric deposition and sequestration of hexachlorocyclohexanes and hexachlorobenzene in the North Atlantic and Arctic oceans. *Environ. Sci. Technol.* 47, 7195–7203.
- Galban-Malagon, C.J., Hernan, G., Abad, E., Dachs, J., 2018. Persistent organic pollutants in krill from the bellingshausen, south scotia, and weddell seas. *Sci. Total Environ.* 610, 1487–1495.
- Gambaro, A., Manodori, L., Zangrando, R., Cincinelli, A., Capodaglio, G., Cescon, P., 2005. Atmospheric PCB concentrations at terra nova bay, Antarctica. *Environ. Sci. Technol.* 39, 9406–9411.
- Garmash, O., Hermanson, M.H., Isaksson, E., Schwikowski, M., Divine, D., Teixeira, C., Muir, D.C.G., 2013. Deposition history of polychlorinated biphenyls to the lomonosovfonna Glacier, Svalbard: A 209 congener analysis. *Environ. Sci. Technol.* 47 (21), 12064–12072.
- Gioia, R., Lohmann, R., Dachs, J., Temme, C., Lakaschus, S., Schulz-Bull, D., Hand, I., Jones, K.C., 2008. Polychlorinated biphenyls in air and water of the North Atlantic and Arctic ocean. *Journal of Geophysical Research-Atmospheres* 113.
- Goerke, H., Weber, K., Bornemann, H., Ramdohr, S., Plotz, J., 2004. Increasing levels and biomagnification of persistent organic pollutants (POPs) in Antarctic biota. *Mar. Pollut. Bull.* 48, 295–302.
- Gong, P., Wang, X., Yao, T., 2011. Ambient distribution of particulate- and gas-phase n-alkanes and polycyclic aromatic hydrocarbons in the Tibetan Plateau. *Environmental Earth Sciences* 64, 1703–1711.
- Gong, P., Wang, X.P., Li, S.H., Yu, W.S., Li, J.L., Kattel, D.B., Wang, W.C., Devkota, L.P., Yao, T.D., Joswiak, D.R., 2014. Atmospheric transport and accumulation of organochlorine compounds on the southern slopes of the Himalayas, Nepal. *Environ. Pollut.* 192, 44–51.
- Gong, P., Wang, X.P., Xue, Y.G., Sheng, J.J., Gao, S.P., Tian, L.D., Yao, T.D., 2015. Influence of atmospheric circulation on the long-range transport of organochlorine pesticides to the western Tibetan Plateau. *Atmos. Res.* 166, 157–164.
- Gouin, T., Mackay, D., Jones, K.C., Harner, T., Meijer, S.N., 2004. Evidence for the "grasshopper" effect and fractionation during long-range atmospheric transport of organic contaminants. *Environ. Pollut.* 128, 139–148.
- Goutte, A., Chevreuil, M., Alliot, F., Chastel, O., Cherel, Y., Eleaume, M., Masse, G., 2013. Persistent organic pollutants in benthic and pelagic organisms off Adélie Land, Antarctica. *Mar. Pollut. Bull.* 77, 82–89.
- Gregor, D.J., Gummer, W.D., 1989. Evidence of atmospheric transport and deposition of organochlorine pesticides and polychlorinated-biphenyls in Canadian arctic snow. *Environ. Sci. Technol.* 23, 561–565.
- Gregor, D.J., Peters, A.J., Teixeira, C., Jones, N., Spencer, C., 1995. The historical residue trend of PCBs in the agassiz ice cap, Ellesmere island, Canada. *Sci. Total Environ.* 160–61, 117–126.
- Grimalt, J.O., Fernandez, P., Berdie, L., Vilanova, R.M., Catalan, J., Psenner, R., Hofer, R., Appleby, P.G., Rosseland, B.O., Lien, L., Massabuau, J.C., Battarbee, R.W., 2001. Selective trapping of organochlorine compounds in mountain lakes of temperate areas. *Environ. Sci. Technol.* 35, 2690–2697.
- Guglielmo, F., Lammel, G., Maier-Reimer, E., 2009. Global environmental cycling of gamma-HCH and DDT in the 1980s-A study using a coupled atmosphere and ocean general circulation model. *Chemosphere* 76, 1509–1517.
- Guo, J., Liao, H., Zhang, L., Zhi, T., Wu, F., 2011. Historical records of sediment polycyclic aromatic hydrocarbons in Qinghai Lake, Northwest China. *Chinese Journal of Ecology* 30, 1467–1472.
- Gustafsson, O., Andersson, P., Axelman, J., Bucheli, T.D., Komp, P., McLachlan, M.S., Sobek, A., Thorngren, J.O., 2005. Observations of the PCB distribution within and in-between ice, snow, ice-rafted debris, ice-interstitial water, and seawater in the Barents Sea marginal ice zone and the North Pole area. *Sci. Total Environ.* 342, 261–279.
- Gustayson, L., Ciesielski, T.M., Bytingsvik, J., Styrisshave, B., Hansen, M., Lie, E., Aars, J., Jenssen, B.M., 2015. Hydroxylated polychlorinated biphenyls decrease circulating steroids in female polar bears (*Ursus maritimus*). *Environ. Res.* 138, 191–201.
- Guzzella, L., Salerno, F., Freppaz, M., Roscioli, C., Pisanello, F., Poma, G., 2016. POP and PAH contamination in the southern slopes of Mt. Everest (Himalaya, Nepal): long-range atmospheric transport, glacier shrinkage, or local impact of tourism? *Sci. Total Environ.* 544, 382–390.
- Halsall, C.J., Barrie, L.A., Fellin, P., Muir, D.C.G., Billeck, B.N., Lockhart, L., Rovinsky, F.Y., Kononov, E.Y., Pastukhov, B., 1997. Spatial and temporal variation of polycyclic aromatic hydrocarbons in the Arctic atmosphere. *Environ. Sci. Technol.* 31, 3593–3599.
- Halsall, C.J., Sweetman, A.J., Barrie, L.A., Jones, K.C., 2001. Modelling the behaviour of PAHs during atmospheric transport from the UK to the Arctic. *Atmos. Environ.* 35, 255–267.
- Hansen, K.M., Christensen, J.H., Brandt, J., Frohn, L.M., Geels, C., 2004. Modelling atmospheric transport of alpha-hexachlorocyclohexane in the Northern Hemisphere with a 3-D dynamical model: DEHM-POP. *Atmos. Chem. Phys.* 4, 1125–1137.
- Hansen, K.M., Christensen, J.H., Geels, C., Silver, J.D., Brandt, J., 2015. Modelling the impact of climate change on the atmospheric transport and the fate of persistent organic pollutants in the Arctic. *Atmos. Chem. Phys.* 15, 6549–6559.
- Harner, T., Kylin, H., Bidleman, T.F., Strachan, W.M.J., 1999. Removal of alpha- and gamma-hexachlorocyclohexane and enantiomers of alpha-hexachlorocyclohexane in the eastern Arctic Ocean. *Environ. Sci. Technol.* 33, 1157–1164.
- Helgason, L.B., Barrett, R., Lie, E., Polder, A., Skaare, J.U., Gabrielsen, G.W., 2008. Levels and temporal trends (1983–2003) of persistent organic pollutants (POPs) and mercury (Hg) in seabird eggs from Northern Norway. *Environ. Pollut.* 155, 190–198.
- Herbert, B.M.J., Halsall, C.J., Villa, S., Jones, K.C., Kallenborn, R., 2005. Rapid changes in PCB and OC pesticide concentrations in Arctic snow. *Environ. Sci. Technol.* 39, 2998–3005.
- Hermanson, M.H., Isaksson, E., Forsstrom, S., Texeira, C., Muir, D.C.G., Pohjola, V.A., van de Wal, R.S.V., 2010. Deposition history of brominated flame retardant compounds in an ice core from Høltedahlfonna, Svalbard, Norway. *Environ. Sci. Technol.* 44, 7405–7410.
- Hermanson, M.H., Isaksson, E.H., Teixeira, C., Muir, D.C.G., Compher, K.M., Li, Y.F., Igarashi, I., Kamiyama, K., 2005. Current-use and legacy pesticide history in the austfonna ice cap, Svalbard, Norway. *Environ. Sci. Technol.* 39, 8163–8169.
- Hoekstra, P.F., O'Hara, T.M., Karlsson, H., Solomon, K.R., Muir, D.C.G., 2003. Enantiomer-specific biomagnification of alpha-hexachlorocyclohexane and selected chiral chlordane-related compounds within an arctic marine food web. *Environ. Toxicol. Chem.* 22, 2482–2491.
- Hong, Q., Wang, Y., Luo, X., Chen, S., Chen, J., Cai, M., Cai, M., Mai, B., 2012. Occurrence of polychlorinated biphenyls (PCBs) together with sediment properties in the surface sediments of the Bering Sea, Chukchi Sea and Canada Basin. *Chemosphere* 88, 1340–1345.
- Hop, H., Borga, K., Gabrielsen, G.W., Kleivane, L., Skaare, J.U., 2002. Food web magnification of persistent organic pollutants in poikilotherms and homeotherms from the Barents Sea. *Environ. Sci. Technol.* 36, 2589–2597.
- Houde, M., De Silva, A.O., Muir, D.C.G., Letcher, R.J., 2011. Monitoring of perfluorinated compounds in aquatic biota: an updated review PFCs in aquatic biota. *Environ. Sci. Technol.* 45, 7962–7973.
- Huang, P., Gong, S.L., Zhao, T.L., Neary, L., Barrie, L.A., 2007. GEM/POPs: a global 3-D dynamic model for semi-volatile persistent organic pollutants - Part 2: global transports and budgets of PCBs. *Atmos. Chem. Phys.* 7, 4015–4025.
- Hung, H., Halsall, C.J., Blanchard, P., Li, H.H., Fellin, P., Stern, G., Rosenberg, B., 2001. Are PCBs in the Canadian Arctic atmosphere declining? Evidence from 5 years of monitoring. *Environ. Sci. Technol.* 35, 1303–1311.
- Hung, H., Katsiyannis, A.A., Brorstrom-Lunden, E., Olafsdottir, K., Aas, W., Breivik, K., Bohlin-Nizzetto, P., Sigurdsson, A., Hakola, H., Bossi, R., Skov, H., Sverko, E., Barresi, E., Fellin, P., Wilson, S., 2016. Temporal trends of persistent organic pollutants (POPs) in arctic air: 20 years of monitoring under the arctic monitoring and assessment Programme (AMAP). *Environ. Pollut.* 217, 52–61.
- Inomata, O.N.K., Montone, R.C., Lara, W.H., Weber, R.R., Toledo, H.H.B., 1996. Tissue distribution of organochlorine residues - PCBs and pesticides - in Antarctic penguins. *Antarct. Sci.* 8, 253–255.
- Jakobsson, M., 2002. Hypsometry and volume of the Arctic Ocean and its constituent seas. *Geochim. Geophys. Geosyst.* 3.
- Jantunen, L.M., Bidleman, T., 1996. Air-water gas exchange of

- hexachlorocyclohexanes (HCHs) and the enantiomers of alpha-HCH in arctic regions. *Journal of Geophysical Research-Atmospheres* 101, 28837–28846.
- Jantunen, L.M., Bidleman, T.F., 1995. Reversal of the air-water gas-exchange direction of hexachlorocyclohexanes in the Bering and Chukchi seas - 1993 versus 1988. *Environ. Sci. Technol.* 29, 1081–1089.
- Jantunen, L.M., Helm, P.A., Kylin, H., Bidleman, T.F., 2008. Hexachlorocyclohexanes (HCHs) in the Canadian archipelago. 2. Air-water gas exchange of alpha- and gamma-HCH. *Environ. Sci. Technol.* 42, 465–470.
- Jantunen, L.M., Wong, F., Gawor, A., Kylin, H., Helm, P.A., Stern, G.A., Strachan, W.M.J., Burniston, D.A., Bidleman, T.F., 2015. 20 Years of air-water gas exchange observations for pesticides in the western Arctic ocean. *Environ. Sci. Technol.* 49, 13844–13852.
- Jantunen, L.M., Bidleman, T.F., 1998. Organochlorine pesticides and enantiomers of chiral pesticides in Arctic Ocean water. *Arch. Environ. Contam. Toxicol.* 35, 218–228.
- Jin, M., Fu, J., Xue, B., Zhou, S., Zhang, L., Li, A., 2017. Distribution and enantiomeric profiles of organochlorine pesticides in surface sediments from the Bering Sea, Chukchi Sea and adjacent Arctic areas. *Environ. Pollut.* 222, 109–117.
- Jin, M., Xue, B., Zhou, S., 2015. Distribution and enantiomeric profiles of organochlorine pesticides in surface sediments from Bering Sea, Chukchi Sea, and Arctic ocean. *Abstr. Pap. Am. Chem. Soc.* 249.
- Jones, K.C., de Voogt, P., 1999. Persistent organic pollutants (POPs): state of the science. *Environ. Pollut.* 100, 209–221.
- Jurado, E., Dachs, J., 2008. Seasonality in the "grasshopping" and atmospheric residence times of persistent organic pollutants over the oceans. *Geophys. Res. Lett.* 35.
- Kallenborn, R., Breivik, K., Eckhardt, S., Lunder, C.R., Mano, S., Schlabach, M., Stohl, A., 2013. Long-term monitoring of persistent organic pollutants (POPs) at the Norwegian Troll station in Dronning Maud Land, Antarctica. *Atmos. Chem. Phys.* 13, 6983–6992.
- Kallenborn, R., Christensen, G., Evensen, A., Schlabach, M., Stohl, A., 2007. Atmospheric transport of persistent organic pollutants (POPs) to Bjørnøya (Bear island). *J. Environ. Monit.* 9, 1082–1091.
- Kallenborn, R., Oehme, M., Wynn-Williams, D.D., Schlabach, M., Harris, J., 1998. Ambient air levels and atmospheric long-range transport of persistent organochlorines to Signy Island, Antarctica. *Sci. Total Environ.* 220, 167–180.
- Kallenborn, R., Reiersen, L.-O., Olseng, C.D., 2012. Long-term atmospheric monitoring of persistent organic pollutants (POPs) in the Arctic: a versatile tool for regulators and environmental science studies. *Atmospheric Pollution Research* 3, 485–493.
- Kang, J.-H., Choi, S.-D., Park, H., Baek, S.-Y., Hong, S., Chang, Y.-S., 2009. Atmospheric deposition of persistent organic pollutants to the east rongbuk glacier in the Himalayas. *Sci. Total Environ.* 408, 57–63.
- Ke, H., Chen, M., Liu, M., Chen, M., Duan, M., Huang, P., Hong, J., Lin, Y., Cheng, S., Wang, X., Huang, M., Cai, M., 2017. Fate of polycyclic aromatic hydrocarbons from the North Pacific to the Arctic: Field measurements and fugacity model simulation. *Chemosphere* 184, 916–923.
- Kelly, B.C., Gobas, F., 2001. Bioaccumulation of persistent organic pollutants in lichen-caribou-wolf food chains of Canada's Central and Western Arctic. *Environ. Sci. Technol.* 35, 325–334.
- Kelly, B.C., Gobas, F., 2003. An arctic terrestrial food-chain bioaccumulation model for persistent organic pollutants. *Environ. Sci. Technol.* 37, 2966–2974.
- Kelly, B.C., Ikonomou, M.G., Blair, J.D., Gobas, F.A.P.C., 2008. Hydroxylated and methoxylated polybrominated diphenyl ethers in a Canadian Arctic marine food web. *Environ. Sci. Technol.* 42, 7069–7077.
- Klanova, J., Matykiewiczova, N., Macka, Z., Prosek, P., Laska, K., Klan, P., 2008. Persistent organic pollutants in soils and sediments from James Ross Island, Antarctica. *Environ. Pollut.* 152, 416–423.
- Krapivin, V.F., 1995. Simulation-model for the investigation of pollution dynamics in the arctic basin. *Okeanologiya* 35, 366–375.
- Kwok, K.Y., Yamazaki, E., Yamashita, N., Taniyasu, S., Murphy, M.B., Horii, Y., Petrick, G., Kallenborn, R., Kannan, K., Murano, K., Lam, P.K.S., 2013. Transport of perfluoroalkyl substances (PFAS) from an arctic glacier to downstream locations: implications for sources. *Sci. Total Environ.* 447, 46–55.
- Lamon, L., von Waldow, H., MacLeod, M., Scheringer, M., Marcomini, A., Hungerbuehler, K., 2009. Modeling the global levels and distribution of polychlorinated biphenyls in air under a climate change scenario. *Environ. Sci. Technol.* 43, 5818–5824.
- Lao, Q., Jiao, L., Chen, F., Chen, L., 2017. Review on researches of legacy POPs and emerging POPs in the arctic regions. *Adv. Earth Sci.* 32, 128–138.
- Lesourd, G.L., Kidd, K.A., De Silva, A.O., Williamson, M., Spencer, C., Wang, X., Muir, D.C.G., 2015. Perfluorinated and polyfluorinated compounds in lake food webs from the Canadian high arctic. *Environ. Sci. Technol.* 49, 2694–2702.
- Letcher, R.J., Bustnes, J.O., Dietz, R., Jenssen, B.M., Jorgensen, E.H., Sonne, C., Verreault, J., Vijayan, M.M., Gabrielsen, G.W., 2010. Exposure and effects assessment of persistent organohalogen contaminants in arctic wildlife and fish. *Sci. Total Environ.* 408, 2995–3043.
- Li, H., Fu, J., Zhang, A., Zhang, Q., Wang, Y., 2016. Occurrence, bioaccumulation and long-range transport of short-chain chlorinated paraffins on the Fildes Peninsula at King George Island, Antarctica. *Environ. Int.* 94, 408–414.
- Li, J., Yuan, G.-L., Li, P., Sun, Y., Yu, H.-H., Wang, G.-H., 2017a. The emerging source of polycyclic aromatic hydrocarbons from mining in the Tibetan Plateau: distributions and contributions in background soils. *Sci. Total Environ.* 584, 64–71.
- Li, J., Yuan, G.-L., Wu, M.-Z., Sun, Y., Han, P., Wang, G.-H., 2017b. Evidence for persistent organic pollutants released from melting glacier in the central Tibetan Plateau, China. *Environ. Pollut.* 220, 178–185.
- Li, J., Yuan, G.-L., Li, P., Sun, Y., Yu, H.-H., Wang, G.-H., 2017c. The emerging source of polycyclic aromatic hydrocarbons from mining in the Tibetan Plateau: distributions and contributions in background soils. *Sci. Total Environ.* 584, 64–71.
- Li, Q., Kang, S., Wang, N., Li, Y., Li, X., Dong, Z., Chen, P., 2017d. Composition and sources of polycyclic aromatic hydrocarbons in cryoconites of the Tibetan Plateau glaciers. *Sci. Total Environ.* 574, 991–999.
- Li, Y., Geng, D., Hu, Y., Wang, P., Zhang, Q., Jiang, G., 2012a. Levels and distribution of polychlorinated biphenyls in the atmosphere close to Chinese Great Wall Station, Antarctica: results from XAD-resin passive air sampling. *Chin. Sci. Bull.* 57, 1499–1503.
- Li, Y., Geng, D., Liu, F., Wang, T., Wang, P., Zhang, Q., Jiang, G., 2012b. Study of PCBs and PBDEs in King George Island, Antarctica, using PUF passive air sampling. *Atmos. Environ.* 51, 140–145.
- Li, Y.F., Macdonald, R.W., 2005. Sources and pathways of selected organochlorine pesticides to the Arctic and the effect of pathway divergence on HCH trends in biota: a review. *Sci. Total Environ.* 342, 87–106.
- Li, Y.F., Macdonald, R.W., Jantunen, L.M., Harner, T., Bidleman, T.F., Strachan, W.M.J., 2002. The transport of beta-hexachlorocyclohexane to the western Arctic Ocean: a contrast to alpha-HCH. *Sci. Total Environ.* 291, 229–246.
- Li, Y.F., Macdonald, R.W., Ma, J.M., Hung, H., Venkatesh, S., 2004. Historical alpha-HCH budget in the Arctic ocean: the arctic mass balance Box model (AMBBM). *Sci. Total Environ.* 324, 115–139.
- Lin, H., Wang, X., Gong, P., Ren, J., Wang, C., Yuan, X., Wang, L., Yao, T., 2017. The influence of climate change on the accumulation of polycyclic aromatic hydrocarbons, black carbon and mercury in a shrinking remote lake of the southern Tibetan Plateau. *Sci. Total Environ.* 601, 1814–1823.
- Liu, F., Liu, Y., Jiang, D., Zhang, R., Cui, Y., Li, M., 2014. Health risk assessment of semi-volatile organic pollutants in Lhasa River China. *Ecotoxicology* 23, 567–576.
- Liu, W., Chen, D., Liu, X., Zheng, X., Yang, W., Westgate, J.N., Wania, F., 2010. Transport of semivolatile organic compounds to the Tibetan plateau: spatial and temporal variation in air concentrations in mountainous western Sichuan, China. *Environ. Sci. Technol.* 44, 1559–1565.
- Lohmann, R., Gioia, R., Jones, K.C., Nizzetto, L., Temme, C., Xie, Z., Schulz-Bull, D., Hand, I., Morgan, E., Jantunen, L., 2009. Organochlorine pesticides and PAHs in the surface water and atmosphere of the North Atlantic and Arctic ocean. *Environ. Sci. Technol.* 43, 5633–5639.
- Lohmann, R., Belkin, I.M., 2014. Organic pollutants and ocean fronts across the Atlantic Ocean. *A Rev.: Prog. Oceanogr.* 128, 172–184.
- Luek, J.L., Dickhut, R.M., Cochran, M.A., Falconer, R.L., Kylin, H., 2017. Persistent organic pollutants in the Atlantic and southern oceans and oceanic atmosphere. *Sci. Total Environ.* 583, 64–71.
- Ma, J., 2010. Atmospheric transport of persistent semi-volatile organic chemicals to the Arctic and cold condensation in the mid-troposphere - Part 1: 2-D modeling in mean atmosphere. *Atmos. Chem. Phys.* 10, 7303–7314.
- Ma, J., Hung, H., Macdonald, R.W., 2016. The influence of global climate change on the environmental fate of persistent organic pollutants: a review with emphasis on the Northern Hemisphere and the Arctic as a receptor. *Glob. Planet. Chang.* 146, 89–108.
- Ma, J., Hung, H., Tian, C., Kallenborn, R., 2011. Revolatilization of persistent organic pollutants in the Arctic induced by climate change. *Nat. Clim. Change* 1, 255–260.
- Ma, J.M., Cao, Z.H., Hung, H., 2004. North Atlantic Oscillation signatures in the atmospheric concentrations of persistent organic pollutants: an analysis using Integrated Atmospheric Deposition Network-Great Lakes monitoring data. *Journal of Geophysical Research-Atmospheres* 109.
- Ma, Y., Halsall, C.J., Crosse, J.D., Graf, C., Cai, M., He, J., Gao, G., Jones, K., 2015. Persistent organic pollutants in ocean sediments from the North Pacific to the Arctic ocean. *Journal of Geophysical Research-Oceans* 120, 2723–2735.
- Ma, Y., Halsall, C.J., Xie, Z., Koetke, D., Mi, W., Ebinghaus, R., Gao, G., 2017a. Polycyclic aromatic hydrocarbons in ocean sediments from the North Pacific to the Arctic ocean. *Environ. Pollut.* 227, 498–504.
- Ma, Y., Xie, Z., Lohmann, R., Mi, W., Gao, G., 2017b. Organophosphate ester flame retardants and plasticizers in ocean sediments from the North Pacific to the Arctic ocean. *Environ. Sci. Technol.* 51, 3809–3815.
- MacLeod, M., Riley, W.J., McKone, T.E., 2005. Assessing the influence of climate variability on atmospheric concentrations of polychlorinated biphenyls using a global-scale mass balance model (BETR-global). *Environ. Sci. Technol.* 39, 6749–6756.
- Malmquist, C., Bindler, R., Renberg, I., Van Bavel, B., Karlsson, E., Anderson, N.J., Tysklind, M., 2003. Time trends of selected persistent organic pollutants in lake sediments from Greenland. *Environ. Sci. Technol.* 37, 4319–4324.
- Mangani, F., Maione, M., Lattanzi, L., Arduini, J., 2001. A ten years monitoring of chlorofluorocarbons at the Antarctica: results and perspectives. *Int. J. Environ. Anal. Chem.* 79, 273–282.
- Martins, C.C., Bicego, M.C., Rose, N.L., Taniguchi, S., Lourenco, R.A., Figueira, R.C.L., Mahiques, M.M., Montone, R.C., 2010. Historical record of polycyclic aromatic hydrocarbons (PAHs) and spheroidal carbonaceous particles (SCPs) in marine sediment cores from Admiralty Bay, King George Island, Antarctica. *Environ. Pollut.* 158, 192–200.
- McDonough, C.A., De Silva, A.O., Sun, C., Cabrerizo, A., Adelman, D., Soltwedel, T., Bauerfeind, E., Dcg, M., Lohmann, R., 2018. Dissolved organophosphate esters and polybrominated diphenyl ethers in remote marine environments: Arctic surface water distributions and net transport through Fram Strait. *Environ. Sci.*

- Technol. 52, 6208–6216.
- McKinney, M.A., Pedro, S., Dietz, R., Sonne, C., Fisk, A.T., Roy, D., Jenssen, B.M., Letcher, R.J., 2015. A review of ecological impacts of global climate change on persistent organic pollutant and mercury pathways and exposures in arctic marine ecosystems. *Current Zoology* 61, 617–628.
- Meijer, S.N., Ockenden, W.A., Sweetman, A., Breivik, K., Grimalt, J.O., Jones, K.C., 2003. Global distribution and budget of PCBs and HCB in background surface soils: implications or sources and environmental processes. *Environ. Sci. Technol.* 37, 667–672.
- Meyer, T., Lei, Y.D., Muradi, I., Wania, F., 2009. Organic contaminant release from melting snow. 2. Influence of snow pack and melt characteristics. *Environ. Sci. Technol.* 43, 663–668.
- Meyer, T., Muir, D.C.G., Teixeira, C., Wang, X., Young, T., Wania, F., 2012. Deposition of brominated flame retardants to the devon ice cap, Nunavut, Canada. *Environ. Sci. Technol.* 46, 826–833.
- Moeller, A., Xie, Z., Sturm, R., Ebinghaus, R., 2010. Large-scale distribution of dechlorane plus in air and seawater from the arctic to Antarctica. *Environ. Sci. Technol.* 44, 8977–8982.
- Moeller, A., Xie, Z., Sturm, R., Ebinghaus, R., 2011. Polybrominated diphenyl ethers (PBDEs) and alternative brominated flame retardants in air and seawater of the European Arctic. *Environ. Pollut.* 159, 1577–1583.
- Moisey, J., Fisk, A.T., Hobson, K.A., Norstrom, R.J., 2001. Hexachlorocyclohexane (HCH) isomers and chiral signatures of alpha-HCH in the arctic marine food web of the Northwest Polynya. *Environ. Sci. Technol.* 35, 1920–1927.
- Montone, R.C., Taniguchi, S., Weber, R.R., 2003. PCBs in the atmosphere of king George island, Antarctica. *Sci. Total Environ.* 308, 167–173.
- Morris, A.D., Muir, D.C.G., Solomon, K.R., Letcher, R.J., McKinney, M.A., Fisk, A.T., McMeans, B.C., Tomy, G.T., Teixeira, C., Wang, X., Duric, M., 2016. Current-use pesticides in seawater and their bioaccumulation in polar bear-ringed seal food chains of the Canadian Arctic. *Environ. Toxicol. Chem.* 35, 1695–1707.
- Morrissey, C.A., Bendall-Young, L.I., Elliott, J.E., 2005. Identifying sources and biomagnification of persistent organic contaminants in biota from mountain streams of southwestern British Columbia, Canada. *Environ. Sci. Technol.* 39, 8090–8098.
- Muir, D., Lohmann, R., 2013. Water as a new matrix for global assessment of hydrophilic POPs. *Trac. Trends Anal. Chem.* 46, 162–172.
- Muir, D.C.G., Norstrom, R.J., Simon, M., 1988. Organochlorine contaminants in Arctic marine-food-chains — accumulation of specific polychlorinated biphenyls and chlordane-related compounds. *Environ. Sci. Technol.* 22, 1071–1079.
- Negoita, T.G., Covaci, A., Gheorghe, A., Schepens, P., 2003. Distribution of polychlorinated biphenyls (PCBs) and organochlorine pesticides in soils from the East Antarctic coast. *J. Environ. Monit.* 5, 281–286.
- Nikanorov, A.M., Bryzgalov, V.A., Reshetnyak, O.S., Kondakova, M.Y., 2015. Pollutant transport in large rivers of the European North and Siberia. *Water Resour.* 42 (3), 322–330.
- Nost, T.H., Halse, A.K., Schlabach, M., Backlund, A., Eckhardt, S., Breivik, K., 2018. Low concentrations of persistent organic pollutants (POPs) in air at Cape Verde. *Sci. Total Environ.* 612, 129–137.
- Octaviani, M., Stemmler, I., Lammel, G., Graf, H.F., 2015. Atmospheric transport of persistent organic pollutants to and from the arctic under present-day and future climate. *Environ. Sci. Technol.* 49, 3593–3602.
- Oehme, M., Mano, S., 1984. The long-rang Transport of organic pollutants to the Arctic. *Fresenius Zeitschrift Fur Analytische Chemie* 319, 141–146.
- Pan, J., Gai, N., Tang, H., Chen, S., Chen, D., Lu, G., Yang, Y., 2014. Organochlorine pesticides and polychlorinated biphenyls in grass, yak muscle, liver, and milk in Ruogai high altitude prairie, the eastern edge of Qinghai-Tibet Plateau. *Sci. Total Environ.* 491–492, 131–137.
- Papale, M., Giannarelli, S., Francesconi, S., Di Marco, G., Mikkonen, A., Conte, A., Rizzo, C., De Domenico, E., Michaud, L., Lo Giudice, A., 2017. Enrichment, isolation and biodegradation potential of psychrotolerant polychlorinated-biphenyl degrading bacteria from the Kongsfjorden (Svalbard Islands, High Arctic Norway). *Mar. Pollut. Bull.* 114, 849–859.
- Park, H., Lee, S.-H., Kim, M., Kim, J.-H., Lim, H.S., 2010. Polychlorinated biphenyl congeners in soils and lichens from king George island, south shetland islands, Antarctica. *Antarct. Sci.* 22, 31–38.
- Pavlova, P.A., Schmid, P., Bogdal, C., Steinlin, C., Jenk, T.M., Schwikowski, M., 2014. Polychlorinated biphenyls in glaciers. 1. deposition history from an alpine ice core. *Environ. Sci. Technol.* 48 (14), 7842–7848.
- Riget, F., Bignert, A., Braune, B., Dam, M., Dietz, R., Evans, M., Green, N., Gunnlaugsdottir, H., Hoydal, K.S., Kucklick, J., Letcher, R., Muir, D., Schuur, S., Sonne, C., Stern, G., Tomy, G., Vorkamp, K., Wilson, S., 2019. Temporal trends of persistent organic pollutants in Arctic marine and freshwater biota. *Sci. Total Environ.* 649, 99–110.
- Pedro, S., Fisk, A.T., Tomy, G.T., Ferguson, S.H., Hussey, N.E., Kessel, S.T., McKinney, M.A., 2017. Mercury and persistent organic pollutants in native and invading forage species of the Canadian Arctic: consequences for food web dynamics. *Environ. Pollut.* 229, 229–240.
- Peters, A.J., Gregor, D.J., Teixeira, C.F., Jones, N.P., Spencer, C., 1995. The recent depositional trend of polycyclic aromatic-hydrocarbons and elemental carbon to the agassiz ice cap, Ellesmere island, Canada. *Sci. Total Environ.* 160–61, 167–179.
- Pokhrel, B., Gong, P., Wang, X., Wang, C., Gao, S., 2018. Polycyclic aromatic hydrocarbons in the urban atmosphere of Nepal: distribution, sources, seasonal trends, and cancer risk. *Sci. Total Environ.* 618, 1583–1590.
- Polkowska, Z., Cichala-Kamrowska, K., Ruman, M., Koziol, K., Krawczyk, W.E., Namiesnik, J., 2011. Organic pollution in surface waters from the fugebekken basin in svalbard, Norwegian arctic. *Sensors* 11, 8910–8929.
- Pozo, K., Martellini, T., Corsolini, S., Harner, T., Estellano, V., Kukucka, P., Mulder, M.A., Lammel, G., Cincinelli, A., 2017. Persistent organic pollutants (POPs) in the atmosphere of coastal areas of the Ross Sea, Antarctica: indications for long-term downward trends. *Chemosphere* 178, 458–465.
- Prevedouros, K., Brorstrom-Lunden, E., Halsall, C.J., Jones, K.C., Lee, R.G.M., Sweetman, A.J., 2004. Seasonal and long-term trends in atmospheric PAH concentrations: evidence and implications. *Environ. Pollut.* 128, 17–27.
- Pucko, M., Stern, G.A., Macdonald, R.W., Barber, D.G., 2010. Alpha- and gamma-hexachlorocyclohexane measurements in the brine fraction of sea ice in the Canadian high arctic using a sump-hole technique. *Environ. Sci. Technol.* 44, 9258–9264.
- Qiu, C., Cai, M., 2010. Ultra trace analysis of 17 organochlorine pesticides in water samples from the Arctic based on the combination of solid-phase extraction and headspace solid-phase microextraction-gas chromatography-electron-capture detector. *J. Chromatogr. A* 1217, 1191–1202.
- Rawn, D.F.K., Lockhart, W.L., Wilkinson, P., Savoie, D.A., Rosenberg, G.B., Muir, D.C.G., 2001. Historical contamination of Yukon Lake sediments by PCBs and organochlorine pesticides: influence of local sources and watershed characteristics. *Sci. Total Environ.* 280, 17–37.
- Ren, J., Wang, X., Wang, C., Gong, P., Wang, X., Yao, T., 2017a. Biomagnification of persistent organic pollutants along a high-altitude aquatic food chain in the Tibetan Plateau: processes and mechanisms. *Environ. Pollut.* 220, 636–643.
- Ren, J., Wang, X.P., Wang, C.F., Gong, P., Yao, T.D., 2017b. Atmospheric processes of organic pollutants over a remote lake on the central Tibetan Plateau: implications for regional cycling. *Atmos. Chem. Phys.* 17, 1401–1415.
- Riget, F., Bignert, A., Braune, B., Stow, J., Wilson, S., 2010a. Temporal trends of legacy POPs in Arctic biota, an update. *Sci. Total Environ.* 408, 2874–2884.
- Riget, F., Vorkamp, K., Bossi, R., Sonne, C., Letcher, R.J., Dietz, R., 2016. Twenty years of monitoring of persistent organic pollutants in Greenland biota. A review. *Environmental Pollution* 217, 114–123.
- Riget, F., Vorkamp, K., Hobson, K.A., Muir, D.C.G., Dietz, R., 2013. Temporal trends of selected POPs and the potential influence of climate variability in a Greenland ringed seal population. *Environmental Science-Processes & Impacts* 15, 1706–1716.
- Riget, F., Vorkamp, K., Muir, D., 2010b. Temporal trends of contaminants in Arctic char (*Salvelinus alpinus*) from a small lake, southwest Greenland during a warming climate. *J. Environ. Monit.* 12, 2252–2258.
- Rios, J.M., Lana, N.B., Ciocco, N.F., Covaci, A., Barrera-Oro, E., Moreira, E., Altamirano, J.C., 2017. Implications of biological factors on accumulation of persistent organic pollutants in Antarctic notothenioid fish. *Ecotoxicol. Environ. Saf.* 145, 630–639.
- Risebrou, R.W., Menzel, D.B., Martin, D.J., Olcott, H.S., 1967. Ddt residues in pacific sea birds — a persistent insecticide in marine food chains. *Nature* 216, 589–8.
- Rudolph, I., Chiang, G., Galban-Malagon, C., Mendoza, R., Martinez, M., Gonzalez, C., Becerra, J., Servos, M.R., Munkittrick, K.R., Barra, R., 2016. Persistent organic pollutants and porphyrins biomarkers in penguin. faeces from Kopaitic Island and Antarctic Peninsula. *Sci. Total Environ.* 573, 1390–1396.
- Ruggirello, R.M., Hermanson, M.H., Isaksson, E., Teixeira, C., Forsstrom, S., Muir, D.C.G., Pohjola, V., van de Wal, R., Meijer, H.A.J., 2010. Current use and legacy pesticide deposition to ice caps on Svalbard, Norway. *J. Geophys. Res.* Atmos. 115.
- Salvado, J.A., Sobek, A., Carrizo, D., Gustafsson, O., 2016. Observation-based assessment of PBDE loads in Arctic ocean waters. *Environ. Sci. Technol.* 50, 2236–2245.
- Sanchis, J., Cabrero, A., Galban-Malagon, C., Barcelo, D., Farre, M., Dachs, J., 2015. Unexpected occurrence of volatile dimethylsiloxanes in antarctic soils. Vegetation, Phytoplankton, and Krill: *Environ. Sci. Technol.* 49, 4415–4424.
- Schiavone, A., Corsolini, S., Borghesi, N., Focardi, S., 2009. Contamination profiles of selected PCB congeners, chlorinated pesticides, PCDD/Fs in Antarctic Fur seal pups and penguin eggs. *Chemosphere* 76, 264–269.
- Schmid, P., Bogdal, C., Bluethgen, N., Anselmetti, F.S., Zwyssig, A., Hungerbuehler, K., 2011. The missing piece: sediment records in remote mountain lakes confirm glaciers being secondary sources of persistent organic pollutants. *Environ. Sci. Technol.* 45, 203–208.
- Sharma, B.M., Nizzetto, L., Bharat, G.K., Tayal, S., Melymuk, L., Sanka, O., Pribylova, P., Audy, O., Larssen, T., 2015. Melting Himalayan glaciers contaminated by legacy atmospheric depositions are important sources of PCBs and high-molecular-weight PAHs for the Ganges floodplain during dry periods. *Environ. Pollut.* 206, 588–596.
- Sheng, J., Wang, X., Gong, P., Joswiak, D.R., Tian, L., Yao, T., Jones, K.C., 2013. Monsoon-driven transport of organochlorine pesticides and polychlorinated biphenyls to the Tibetan plateau: three year atmospheric monitoring study. *Environ. Sci. Technol.* 47, 3199–3208.
- Sladen, W.J.L., Menzie, C.M., Reichel, W.L., 1966. Ddt residues in adielie penguins and a crabeater seal from Antarctica. *Nature* 210, 670–8.
- Sobek, A., Gustafsson, O., 2004. Latitudinal fractionation of polychlorinated biphenyls in surface seawater along a 62 degrees N–89 degrees N transect from the southern Norwegian Sea to the North Pole area. *Environ. Sci. Technol.* 38, 2746–2751.
- Sobek, A., Gustafsson, O., 2014a. Deep water masses and sediments are main compartments for polychlorinated biphenyls in the Arctic ocean. *Environ. Sci. Technol.* 48, 6719–6725.
- Sobek, A., Gustafsson, O., 2014b. Deep water masses and sediments are main

- compartments for polychlorinated biphenyls in the Arctic ocean. *Environ. Sci. Technol.* 48, 6719–6725.
- Stemmler, I., Lammel, G., 2009. Cycling of DDT in the global environment 1950–2002: world ocean returns the pollutant. *Geophys. Res. Lett.* 36.
- Stemmler, I., Lammel, G., 2010. Pathways of PFOA to the Arctic: variabilities and contributions of oceanic currents and atmospheric transport and chemistry sources. *Atmos. Chem. Phys.* 10, 9965–9980.
- Stemmler, I., Lammel, G., 2013. Evidence of the return of past pollution in the ocean: a model study. *Geophys. Res. Lett.* 40.
- Stow, J.P., Sova, J., Reimer, K.J., 2005. The relative influence of distant and local (DEW-line) PCB sources in the Canadian Arctic. *Sci. Total Environ.* 342, 107–118.
- Sun, C., Soltwedel, T., Bauerfeind, E., Adelman, D.A., Lohmann, R., 2016. Depth profiles of persistent organic pollutants in the North and tropical Atlantic ocean. *Environ. Sci. Technol.* 50, 6172–6179.
- Sun, L.G., Yin, X.B., Pan, C.P., Wang, Y.H., 2005. A 50-years record of dichlorodiphenyl-trichloroethanes and hexachlorocyclohexanes in lake sediments and penguin droppings on King George Island, Maritime Antarctic. *J. Environ. Sci.* 17, 899–905.
- Sun, Y., Yuan, G.L., Li, J., Tang, J.H., Wang, G.H., 2018. High-resolution sedimentary records of some organochlorine pesticides in Yamzho Yumco Lake of the Tibetan Plateau: concentration and composition. *Sci. Total Environ.* 615, 469–475.
- Tao, S., Wang, W., Liu, W., Zuo, Q., Wang, X., Wang, R., Wang, B., Shen, G., Yang, Y., He, J.-S., 2011. Polycyclic aromatic hydrocarbons and organochlorine pesticides in surface soils from the Qinghai-Tibetan plateau. *J. Environ. Monit.* 13, 175–181.
- Tao, Y.Q., Lei, G.L., Xue, B., Yao, S.C., Pu, Y., Zhang, H.C., 2014. Deposition and regional distribution of HCHs and p,p'-DDX in the western and southern Tibetan Plateau: records from a lake sediment core and the surface soils. *Environ. Sci. Pollut. Control Ser.* 21, 1875–1883.
- Uhl, S., Scheringer, M., Stohl, A., Burkhardt, J.F., Hungerbühler, K., 2012. Primary source regions of polychlorinated biphenyls (PCBs) measured in the Arctic. *Atmos. Environ.* 62, 391–399.
- UNEP/AMAP, 2011. Climate Change and POPs: Predicting the Impacts. Report of the UNEP/AMAP Expert Group. Secretariat of the Stockholm Convention, Geneva, p. 62.
- Valera, B., Ayotte, P., Poirier, P., Dewailly, E., 2013. Associations between plasma persistent organic pollutant levels and blood pressure in Inuit adults from Nunavik. *Environ. Int.* 59, 282–289.
- van den Brink, N.W., Riddle, M.J., van den Heuvel-Greve, M., van Franeker, J.A., 2011. Contrasting time trends of organic contaminants in Antarctic pelagic and benthic food webs. *Mar. Pollut. Bull.* 62, 128–132.
- van Drooge, B.L., Grimalt, J.O., Stuchlik, E., 2013. Spatial distribution of polychlorinated biphenyls in High Tatra lake sediments. *Environ. Sci. Pollut. Control Ser.* 20, 6594–6600.
- Vecchiato, M., Argiriadis, E., Zambon, S., Barbante, C., Toscano, G., Gambaro, A., Piazza, R., 2015. Persistent Organic Pollutants (POPs) in Antarctica: occurrence in continental and coastal surface snow. *Microchem. J.* 119, 75–82.
- Veilleux, J., Muir, D.C.G., Antoniadou, D., Small, J.M., Spencer, C., Loewen, T.N., Babaluk, J.A., Reist, J.D., Vincent, W.F., 2012. Perfluorinated chemicals in meromictic lakes on the northern coast of Ellesmere island, high arctic Canada. *Arctic* 65, 245–256.
- Villa, S., Migliorati, S., Monti, G.S., Holoubek, I., Vighi, M., 2017. Risk of POP mixtures on the arctic food chain. *Environ. Toxicol. Chem.* 36, 1181–1192.
- Vorkamp, K., 2016. An overlooked environmental issue? A review of the inadvertent formation of PCB-11 and other PCB congeners and their occurrence in consumer products and in the environment. *Sci. Total Environ.* 541, 1463–1476.
- Vorkamp, K., Riget, F.F., 2014. A review of new and current-use contaminants in the Arctic environment: evidence of long-range transport and indications of bioaccumulation. *Chemosphere* 111, 379–395.
- Wang, C.F., Gong, P., Wang, X., Yao, T., 2016a. Distribution, Environmental Behavior, and health risks of polychlorinated biphenyls in the Tibetan agricultural soil and crops. *Asian J. Ecotoxicol.* 11, 339–346.
- Wang, C.F., Wang, X.P., Gong, P., Yao, T., 2014a. Polycyclic aromatic hydrocarbons in surface soil across the Tibetan Plateau: spatial distribution, source and air-soil exchange. *Environ. Pollut.* 184, 138–144.
- Wang, C.F., Wang, X.P., Gong, P., Yao, T., 2016b. Residues, spatial distribution and risk assessment of DDTs and HCHs in agricultural soil and crops from the Tibetan Plateau. *Chemosphere* 149, 358–365.
- Wang, C.F., Wang, X.P., Gong, P., Yao, T., 2017a. Long-term trends of atmospheric organochlorine pollutants and polycyclic aromatic hydrocarbons over the southeastern Tibetan Plateau. *Sci. Total Environ.* 624, 241–249.
- Wang, C.F., Wang, X.P., Yuan, X., Ren, J., Gong, P., 2015a. Organochlorine pesticides and polychlorinated biphenyls in air, grass and yak butter from Namco in the central Tibetan Plateau. *Environ. Pollut.* 201, 50–57.
- Wang, C.F., Wang, X.P., Gong, P., Yao, T.D., 2014b. Polycyclic aromatic hydrocarbons in surface soil across the Tibetan Plateau: spatial distribution, source and air-soil exchange. *Environ. Pollut.* 184, 138–144.
- Wang, P., Meng, W.Y., Li, Y.M., Zhang, Q.H., Zhang, L., Fu, J., Yang, R.Q., Jiang, G.B., 2018. Temporal variation (2011–2014) of atmospheric OCPs at king George island, west Antarctica. *Atmos. Environ.* 191, 432–439.
- Wang, P., Zhang, Q., Yang, Q., Zhang, L., Liu, F., Fu, J., Meng, W., Wang, D., Sun, H., Zheng, S., Hao, Y., Liang, Y., Jiang, G.B., 2017b. Three-year monitoring of atmospheric PCBs and PBDEs at the Chinese Great Wall Station, West Antarctica: levels, chiral signature, environmental behaviors and source implication. *Atmos. Environ.* 150, 407–416.
- Wang, P., Zhang, Q.-h., Thanh, W., Chen, W.-h., Ren, D.-w., Li, Y.-m., Jiang, G.-b., 2012a. PCBs and PBDEs in environmental samples from king George island and ardley island, Antarctica. *RSC Adv.* 2, 1350–1355.
- Wang, P., Zhang, Q.H., Wang, Y.W., Wang, T., Li, X.M., Li, Y.M., Ding, L., Jiang, G.B., 2009a. Altitude dependence of polychlorinated biphenyls (PCBs) and polybrominated diphenyl ethers (PBDEs) in surface soil from Tibetan Plateau, China. *Chemosphere* 76, 1498–1504.
- Wang, Q., Shi, Y., Hu, J., Yao, Z., Fang, X., Dong, Y., 2012b. Determination of dioxin-like polychlorinated biphenyls in soil and moss from Fildes Peninsula, Antarctica. *Chin. Sci. Bull.* 57, 992–996.
- Wang, S., Ni, H.G., Sun, J.L., Jing, X., He, J.S., Zeng, H., 2013. Polycyclic aromatic hydrocarbons in soils from the Tibetan Plateau, China: distribution and influence of environmental factors. *Environmental Science-Processes & Impacts* 15, 661–667.
- Wang, X.P., Xu, B.Q., Kang, S.C., Cong, Z.Y., Yao, T.D., 2008a. The historical residue trends of DDT, hexachlorocyclohexanes and polycyclic aromatic hydrocarbons in an ice core from Mt. Everest, central Himalayas, China. *Atmos. Environ.* 42, 6699–6709.
- Wang, X.P., Gong, P., Yao, T.-D., Jones, K.C., 2010a. Passive air sampling of organochlorine pesticides, polychlorinated biphenyls, and polybrominated diphenyl ethers across the Tibetan plateau. *Environ. Sci. Technol.* 44, 2988–2993.
- Wang, X.P., Gong, P., Wang, C., Ren, J., Yao, T., 2016c. A review of current knowledge and future prospects regarding persistent organic pollutants over the Tibetan Plateau. *Sci. Total Environ.* 573, 139–154.
- Wang, X.P., Halsall, C., Codling, G., Xie, Z., Xu, B., Zhao, Z., Xue, Y., Ebinghaus, R., Jones, K.C., 2014c. Accumulation of perfluoroalkyl compounds in Tibetan mountain snow: temporal patterns from 1980 to 2010. *Environ. Sci. Technol.* 48, 173–181.
- Wang, X.P., Ren, J., Gong, P., Wang, C., Xue, Y., Yao, T., Lohmann, R., 2016d. Spatial distribution of the persistent organic pollutants across the Tibetan Plateau and its linkage with the climate systems: a 5-year air monitoring study. *Atmos. Chem. Phys.* 16, 6901–6911.
- Wang, X., Xue, Y., Gong, P., Yao, T., 2014d. Organochlorine pesticides and polychlorinated biphenyls in Tibetan forest soil: profile distribution and processes. *Environ. Sci. Pollut. Control Ser.* 21, 1897–1904.
- Wang, X.P., Yang, H., Gong, P., Zhao, X., Wu, G., Turner, S., Yao, T., 2010b. One century sedimentary records of polycyclic aromatic hydrocarbons, mercury and trace elements in the Qinghai Lake, Tibetan Plateau. *Environ. Pollut.* 158, 3065–3070.
- Wang, X.P., Sheng, J.J., Gong, P., Xue, Y.G., Yao, T.D., Jones, K.C., 2012c. Persistent organic pollutants in the Tibetan surface soil: spatial distribution, air-soil exchange and implications for global cycling. *Environ. Pollut.* 170, 145–151.
- Wang, X.P., Yao, T.D., Wang, P.L., Yang, W., Tian, L.D., 2008b. The recent deposition of persistent organic pollutants and mercury to the Dasuopu glacier, Mt. Xixiabangma, central Himalayas. *Sci. Total Environ.* 394, 134–143.
- Wang, Z., Ma, X.D., Na, G.S., Lin, Z.S., Ding, Q., Yao, Z.W., 2009b. Correlations between physicochemical properties of PAHs and their distribution in soil, moss and reindeer dung at Ny-Alesund of the Arctic. *Environ. Pollut.* 157, 3132–3136.
- Wang, Z., Na, G.S., Ma, X.D., Ge, L.K., Lin, Z.S., Yao, Z.W., 2015b. Characterizing the distribution of selected PBDEs in soil, moss and reindeer dung at Ny-Alesund of the Arctic. *Chemosphere* 137, 9–13.
- Wania, F., Binnington, M.J., Curren, M.S., 2017. Mechanistic modeling of persistent organic pollutant exposure among indigenous Arctic populations: motivations, challenges, and benefits. *Environ. Rev.* 25, 396–407.
- Wania, F., Dugani, C.B., 2003. Assessing the long-range transport potential of polybrominated diphenyl ethers: a comparison of four multimedia models. *Environ. Toxicol. Chem.* 22, 1252–1261.
- Wania, F., Mackay, D., 1993. Global fractionation and cold condensation of low volatility organochlorine compounds in polar regions. *Ambio* 2, 10–18.
- Wania, F., Mackay, D., 1995. A global distribution model for persistent organic chemicals. *Sci. Total Environ.* 160–61, 211–232.
- Wania, F., Mackay, D., 1999. Global chemical fate of alpha-hexachlorocyclohexane. 2. Use of a global distribution model for mass balancing, source apportionment, and trend prediction. *Environ. Toxicol. Chem.* 18, 1400–1407.
- Wania, F., Mackay, D., Li, Y.F., Bidleman, T.F., Strand, A., 1999. Global chemical fate of alpha-hexachlorocyclohexane. 1. Evaluation of a global distribution model. *Environ. Toxicol. Chem.* 18, 1390–1399.
- Wania, F., Westgate, J.N., 2008. On the mechanism of mountain cold-trapping of organic chemicals. *Environ. Sci. Technol.* 42, 9092–9098.
- Weber, J., Halsall, C.J., Muir, D.C.G., Teixeira, C., Burniston, D.A., Strachan, W.M.J., Hung, H., Mackay, N., Arnold, D., Kylin, H., 2006. Endosulfan and gamma-HCH in the Arctic: an assessment of surface seawater concentrations and air-sea exchange. *Environ. Sci. Technol.* 40, 7570–7576.
- Welch, H.E., Muir, D.C.G., Billeck, B.N., Lockhart, W.L., Brunskill, G.J., Kling, H.J., Olson, M.P., Lemoine, R.M., 1991. Brown snow – a long-range transport event in the Canadian Arctic. *Environ. Sci. Technol.* 25, 280–286.
- Woehrschimmel, H., MacLeod, M., Hungerbühler, K., 2013. Emissions, fate and transport of persistent organic pollutants to the arctic in a changing global climate. *Environ. Sci. Technol.* 47, 2323–2330.
- Wong, F., Jantunen, L.M., Pucko, M., Papakyriakou, T., Staebler, R.M., Stern, G.A., Bidleman, T.F., 2011. Air-water exchange of anthropogenic and natural organohalogenes on international polar year (IPY) expeditions in the Canadian arctic. *Environ. Sci. Technol.* 45, 876–881.
- Wong, F., Shoen, M., Katsoyiannis, A., Eckhardt, S., Stohl, A., Bohlin-Nizzetto, P., Li, H., Fellin, P., Su, Y., Hung, H., 2018. Assessing temporal trends and source regions of per- and polyfluoroalkyl substances (PFASs) in air under the Arctic Monitoring and Assessment Programme (AMAP). *Atmos. Environ.* 172, 65–73.

- Xie, Z., Koch, B.P., Moeller, A., Sturm, R., Ebinghaus, R., 2011. Transport and fate of hexachlorocyclohexanes in the oceanic air and surface seawater. *Biogeosciences* 8, 2621–2633.
- Xue, R., Chen, L., Lu, Z., Wang, J., Yang, H., Zhang, J., Cai, M., 2016. Spatial distribution and source apportionment of PAHs in marine surface sediments of Prydz Bay, East Antarctica. *Environ. Pollut.* 219, 528–536.
- Yamashita, N., Kannan, K., Taniyasu, S., Horii, Y., Petrick, G., Gamo, T., 2005. A global survey of perfluorinated acids in oceans. *Mar. Pollut. Bull.* 51, 658–668.
- Yamashita, N., Taniyasu, S., Petrick, G., Wei, S., Gamo, T., Lam, P.K.S., Kannan, K., 2008. Perfluorinated acids as novel chemical tracers of global circulation of ocean waters. *Chemosphere* 70, 1247–1255.
- Yang, R., Jing, C., Zhang, Q., Wang, Z., Wang, Y., Li, Y., Jiang, G., 2011. Polybrominated diphenyl ethers (PBDEs) and mercury in fish from lakes of the Tibetan Plateau. *Chemosphere* 83, 862–867.
- Yang, R., Wang, Y., Li, A., Zhang, Q., Jing, C., Wang, T., Wang, P., Li, Y., Jiang, G., 2010. Organochlorine pesticides and PCBs in fish from lakes of the Tibetan Plateau and the implications. *Environ. Pollut.* 158, 2310–2316.
- Yang, R., Xie, T., Li, A., Yang, H., Turner, S., Wu, G., Jing, C., 2016. Sedimentary records of polycyclic aromatic hydrocarbons (PAHs) in remote lakes across the Tibetan Plateau. *Environ. Pollut.* 214, 1–7.
- Yang, R., Zhang, S., Li, A., Jiang, G., Jing, C., 2013. Altitudinal and spatial signature of persistent organic pollutants in soil, lichen, conifer needles, and bark of the southeast Tibetan plateau: implications for sources and environmental cycling. *Environ. Sci. Technol.* 47, 12736–12743.
- Yao, T.D., Masson-Delmotte, V., Gao, J., Yu, W.S., Yang, X.X., Risi, C., Sturm, C., Werner, M., Zhao, H.B., He, Y., Ren, W., Tian, L.D., Shi, C.M., Hou, S.G., 2013. A review of climate controls on delta O-18 in precipitation over the Tibetan Plateau: observations and simulations. *Rev. Geophys.* 51.
- Yao, Z.W., Jiang, G.B., Cai, Y.Q., Xu, H.Z., Ma, Y.A., 2003. Status of persistent organic pollutants and heavy metals in surface water of Arctic region. *Chin. Sci. Bull.* 48, 131–135.
- Yao, Z.W., Jiang, G.B., Xu, H.Z., 2002. Distribution of organochlorine pesticides in seawater of the bering and Chukchi Sea. *Environ. Pollut.* 116, 49–56.
- Yeung, L.W.Y., Dassuncao, C., Mabury, S., Sunderland, E.M., Zhang, X., Lohmann, R., 2017. Vertical profiles, sources, and transport of PFASs in the Arctic ocean. *Environ. Sci. Technol.* 51, 6735–6744.
- Yogui, G.T., Sericano, J., 2008. Polybrominated diphenyl ether flame retardants in lichens and mosses from King George Island, maritime Antarctica. *Chemosphere* 73, 1589–1593.
- Yogui, G.T., Sericano, J.L., Montone, R.C., 2011. Accumulation of semivolatile organic compounds in Antarctic vegetation: a case study of polybrominated diphenyl ethers. *Sci. Total Environ.* 409, 3902–3908.
- Young, C.J., Furdui, V., Franklin, J., Koerner, R.M., Muir, D.C.G., Mabury, S.A., 2007. Perfluorinated acids in Arctic snow: new evidence for atmospheric formation. *Environ. Sci. Technol.* 41 (10), 3455–3461.
- Zaborska, A., Carroll, J., Pazdro, K., Pempkowiak, J., 2011. Spatio-temporal patterns of PAHs, PCBs and HCB in sediments of the western Barents Sea. *Oceanologia* 53, 1005–1026.
- Zhang, L., Ma, J., Tian, C., Li, Y., Hung, H., 2010. Atmospheric transport of persistent semi-volatile organic chemicals to the Arctic and cold condensation in the mid-troposphere - Part 2: 3-D modeling of episodic atmospheric transport. *Atmos. Chem. Phys.* 10, 7315–7324.
- Zhang, X., Meyer, T., Muir, D.C.G., Teixeira, C., Wang, X., Wania, F., 2013. Atmospheric deposition of current use pesticides in the arctic: snow core records from the Devon island ice cap, nunavut, Canada. *Environmental Science-Processes & Impacts* 15, 2304–2311.
- Zhang, X., Zhang, Y., Dassuncao, C., Lohmann, R., Sunderland, E.M., 2017. North Atlantic Deep Water formation inhibits high Arctic contamination by continental perfluorooctane sulfonate discharges. *Glob. Biogeochem. Cycles* 31, 1332–1343.
- Zhao, Z., Xie, Z., Moeller, A., Sturm, R., Tang, J., Zhang, G., Ebinghaus, R., 2012. Distribution and long-range transport of polyfluoroalkyl substances in the Arctic, Atlantic Ocean and Antarctic coast. *Environ. Pollut.* 170, 71–77.
- Zheng, X., Liu, X., Jiang, G., Wang, Y., Zhang, Q., Cai, Y., Cong, Z., 2012. Distribution of PCBs and PBDEs in soils along the altitudinal gradients of Balang Mountain, the east edge of the Tibetan Plateau. *Environ. Pollut.* 161, 101–106.
- Zhou, R.C., Yang, R.Q., Jing, C.Y., 2018. Polycyclic aromatic hydrocarbons in soils and lichen from the western Tibetan Plateau: concentration profiles, distribution and its influencing factors. *Ecotoxicol. Environ. Saf.* 152, 151–158.
- Zhu, C., Li, Y., Wang, P., Chen, Z., Ren, D., Ssebugere, P., Zhang, Q., Jiang, G., 2015a. Polychlorinated biphenyls (PCBs) and polybrominated biphenyl ethers (PBDEs) in environmental samples from Ny-Alesund and London Island, Svalbard, the Arctic. *Chemosphere* 126, 40–46.
- Zhu, N., Fu, J., Gao, Y., Ssebugere, P., Wang, Y., Jiang, G., 2013. Hexabromocyclododecane in alpine fish from the Tibetan Plateau, China. *Environ. Pollut.* 181, 7–13.
- Zhu, N., Schramm, K.-W., Wang, T., Henkelmann, B., Fu, J., Gao, Y., Wang, Y., Jiang, G., 2015b. Lichen, moss and soil in resolving the occurrence of semi-volatile organic compounds on the southeastern Tibetan Plateau, China. *Sci. Total Environ.* 518, 328–336.
- Zhulidov, A.V., Headley, J.V., Pavlov, D.F., Robarts, R.D., Korotova, L.G., Vinnikov, Y.Y., Zhulidova, O.V., 2000. Riverine fluxes of the persistent organochlorine pesticides hexachlorocyclohexane and DDT in the Russian Federation. *Chemosphere* 41, 829–841.
- Zhulidov, A.V., Robarts, R.D., Headley, J.V., Liber, K., Zhulidov, D.A., Zhulidova, O.V., Pavlov, D.F., 2002. Levels of DDT and hexachlorocyclohexane in burbot (*Lota lota* L.) from Russian Arctic rivers. *Sci. Total Environ.* 292, 231–246.